100 PM

E. Reggij

امتحانات رقورا)







Cairo Governorate



Shoubra Educational Zone **Mathematics Supervision**

First

Multiple choice questions



Cho

ose the correct answer from the given ones:	Interactive test (1)
If $7^{x} = 4$, then $x = \dots$	

- (1)
 - (a) $\frac{4}{7}$

- (b) $\frac{7}{4}$
- $(c) \log_7 4$
- $(d) \log_4 7$
- (2) The point of symmetry of the curve of the function $f: f(X) = X^3$ is
 - (a)(1,1)
- (b) (0,0)
- (c)(1,0)
- (d)(0,1)

- (3) If $2^{x-5} = 3^{5-x}$, then $x = \dots$
 - (a) $\frac{2}{3}$

- (b) $\frac{3}{2}$
- (c)0
- (d)5
- (4) The curve of the function g: g(x) = |x + 3| is the same curve of the function f: f(X) = |X| by a translation 3 units in the direction of
 - (a) \overrightarrow{OX}

- (b) OX
- (c) \overrightarrow{OY}
- (d) OY

- (5) If $f(X) = 5^X$, then $\frac{f(X+2)}{f(X+1)} = \dots$
 - (a) 25

- (b) 5
- (c) 1
- (d) $\frac{1}{5}$

- (6) $\lim_{x \to 1} \frac{x^2 1}{x^2 + 1} = \dots$
 - (a) 0

- (c) 2
- (d) doesn't exist.

- (7) $\lim_{x \to 3} \frac{x^2 8x + 15}{x 3} = \dots$

- (b) 3
- (c) 2
- (d) 2
- (8) The measure of the greatest angle of \triangle ABC where : a = 3 cm., b = 4 cm., c = 5 cm. equals
 - (a) 90°

- (b) 60°
- (c) 30°
- (d) 120°

- (9) In \triangle ABC: if a = 7 cm., then $b = \cdots$
 - (a) $\frac{\sin A}{7 \sin B}$

- (b) $\frac{\sin C}{7 \sin B}$ (c) $\frac{7 \sin B}{\sin A}$
- $\frac{\text{(d)}}{\sin A}$
- (10) If f is an even function on the interval $a \cdot b$, then $b = \cdots$
 - (a) a

- (b) a
- (c) 2 a
- $(d) a^2$

(11) The domain of the function $f(x) = \frac{x-3}{x^2-5}$ is

(a) R

(b) $\mathbb{R} - \{3\}$ (c) $\mathbb{R} - \{-2, -3\}$ (d) $\mathbb{R} - \{2, 3\}$

(12) If $f(x) = 2^x$, then $f(-1) = \cdots$

(a) 1

(b) - 1

(c) $\frac{1}{2}$

(d) $\frac{-1}{2}$

(13) If $\left(\frac{1}{2}\right)^{x} = 8$, then $x = \dots$

(b) 3

(c) $\frac{1}{2}$

 $(d) \frac{-1}{2}$

(14) $\lim_{x \to 1} \frac{x^5 - 1}{x - 1} = \dots$

(b) 5

(c) 4

(d) 1

(15) $\lim_{x \to 4} \frac{x-4}{\sqrt{5+x}-3} = \dots$

(c) 6

(d) 3

(16) If $\lim_{x \to 3} \frac{a}{x+1} = 2$, then $a = \dots$

(a) 2

(b) 4

(c) 6

(d) 8

(17) The radius length of the circumcircle of \triangle ABC in which: m (\angle A) = 30°, a = 10 cm. equals cm.

(a) 5

(b) 10

(c) 20

(d) 40

(18) In \triangle ABC if $a^2 = b^2 + c^2 + bc$, then m (\triangle A) =

(c) 120°

(d) 150°

(19) $\log 25 + \frac{\log 8 \times \log 16}{\log 64} = \cdots$

(a) 2

(c) 10

(d) 100

(20) The solution set of the equation : $|2 \times -1| = 5$ in \mathbb{R} is

(a) R

(b) [-2,3] (c) $\{3\}$

 $(d) \{-2,3\}$

(21) If $\log 3 = X$, $\log 4 = y$, then $\log 12 = \dots$

(a) $\log x + \log y$

(b) $\chi - v$

(c) $\chi + y$

 $(d) \chi y$

(22) If $\log_2 x = \frac{1}{3}$, then $\log_2 (8 x^3) = \dots$

(a) 1

(c) 2

(d)3

(23) $\lim_{x \to \infty} \left(3 - \frac{7}{x} + \frac{4}{x^2} \right) = \dots$

(b) 0

(c) - 7

(d) 3

(24) $\lim_{x \to 7} \frac{x^3 - 343}{x^2 - 49} = \cdots$

(a) $\frac{21}{2}$

(b) $\frac{2}{21}$

(c) $\frac{3}{2}$

(d) 14

(25)
$$\lim_{x \to \infty} \frac{2x^2 + 5x - 3}{7 - 3x^2} = \dots$$

(a) $\frac{-3}{7}$ (b) $\frac{-2}{3}$

(a)
$$\frac{-3}{7}$$

(b)
$$\frac{-2}{3}$$

(26) In \triangle ABC: if a = 3 cm., b = 4 cm., c = 6 cm., then $\cos C = \cdots$

(a)
$$\frac{-11}{24}$$

(b)
$$\frac{11}{24}$$

(c)
$$\frac{-11}{12}$$
 (d) $\frac{11}{12}$

(d)
$$\frac{11}{12}$$

(27) If r is the radius length of the circumcircle of \triangle ABC, then $\frac{2 \text{ b}}{\sin \text{ B}} = \cdots$

(a)
$$\frac{1}{2}$$
 r

(b) r

(c) 2 r

(d) 4 r

Second Essay questions

Answer the following questions :

If Graph the function $f(x) = x^2 - 3$, then from the graph deduce the range of the function , its monotony and its type whether it is even , odd or otherwise.

2 Find:
$$\lim_{x \to \infty} \frac{16 x^{-4} - 7 x^{-1} - 27}{8 x^{-4} - 9}$$

Cairo Governorate



Zeiton Educational Zone

Multiple choice questions First



Choose the correct answer from the given ones:

Interactive

(1) The domain of the function $f(x) = \frac{7}{x^3 - x}$ is

(a)
$$\mathbb{R} - \{3\}$$

(b)
$$\mathbb{R} - \{7\}$$

(c)
$$\mathbb{R} - \{0, 1\}$$

(b)
$$\mathbb{R} - \{7\}$$
 (c) $\mathbb{R} - \{0, 1\}$ (d) $\mathbb{R} - \{0, -1, 1\}$

(2) All the following are unspecified quantities except

$$(c) \infty + \infty$$
 $(d) \infty \div \infty$

(3) The exponential function of base "a" is increasing function if

(a)
$$a > 0$$

(b)
$$a > 1$$

(c)
$$0 < a < 1$$

(d)
$$a = 1$$

(4) $\lim_{x \to 3} \frac{2x-6}{7x-21} = \cdots$

(a)
$$\frac{2}{3}$$

(b)
$$\frac{2}{7}$$

(c)
$$\frac{3}{7}$$

$$(d)$$
 3

(5) XYZ is an equilateral triangle the length of its side is $10\sqrt{3}$ cm., then the length of the diameter of its circumcircle is cm.

(6) If $2^{X+1} = 5^{X+1}$, then $3^{X+1} = \dots$

$$(c) - 1$$

$$(d)$$
 3

$$\begin{array}{cc} (7) \lim_{x \to \infty} \frac{\sqrt{x^2}}{x} = \cdots \end{array}$$

(b) 1

(c) 2

(d) - 1

(8) If $\log_3 X = 2$, then $X = \cdots$

(c) 8

(d) 9

(b) 5 $\lim_{x \to 4} (3 x - \sqrt{x}) = \dots$ (a) 8

(c) 14

(d) 16

(10) The domain of the function $f: f(x) = \sqrt{x-3}$ is

(b) $\mathbb{R} - \{3\}$

(c) [3,∞[

 $(d) - \infty, 3$

(11) If "r" is the length of the radius of the circumcircle of the triangle XYZ

, then
$$\frac{y}{2 \sin Y} = \cdots$$

(b) 2 r

(c) $\frac{1}{2}$ r

(d) 4 r

(12) If $\log_3 (2 X + 3) = 2$, then $X = \dots$

(c) 9

(d) 4

(13) $\lim_{x \to 0} \frac{7 + 2x}{\cos x} = \dots$

(b) 8

(c) 9

(d) 1

(14) The function $f: f(X) = X \cos X$

(a) even.

(b) odd.

(c) neither even nor odd.

(d) linear.

(15) $\lim_{x \to 2} \frac{x^5 - 32}{x^3 - 8} = \dots$

(a) 4

(b) $\frac{5}{3}$

(d) $6\frac{2}{3}$

(16) If \angle A supplement of \angle C, then \cos A + \cos C =

(a) zero

(b) 1

(d) $\frac{1}{2}$

(17) If $2^{x} = 4^{y} = 64$, then $x + y = \dots$

(d) 9

(18) $\lim_{x \to 0} \frac{\sqrt[3]{x+1}-1}{x} = \dots$

(a) 1

(c) zero

 $(d) - \frac{2}{3}$

(19) The solution set of $\sqrt{4 x^2 - 12 x + 9} \le 9$ is

(a) [-6, 12]

(b) [-3,6] (c) $\mathbb{R} - [-3,6]$ (d) $\mathbb{R} -]-3,6[$

(20) In \triangle XYZ: $y^2 + z^2 - x^2 = 2$ y z ×

(a) cos X

(b) sin Z

(c) $\cos Z$ (d) $\sin X$

(21) $\log_2 5 \times \log_5 2 = \dots$

(b) 10

(c) $\log_2 10$ (d) $\log_5 10$

(22) $\lim_{x \to \infty} \left(\frac{3}{x^2} - 2 \right) = \cdots$

(b) 2

(c) - 3

(23) The symmetric point of the curve of the function $f: f(x) = \frac{1}{x-3} + 4$ is

(a) (3, -4)

(b) (-3, -4)

(c)(3,4)

(24) In triangle ABC if a = 5 cm., b = 7 cm. and $m (\angle C) = 65^{\circ}$, then $c = \cdots cm$.

(a) 44.4

(b) 32.1

(c) 6.7

(d) 8.2

 $(25) \frac{1}{\log_2 14} + \frac{1}{\log_7 14} = \dots$

(c) 7

(d) 14

(26) If the function $f: f(X) = a^X$ passing through the point (1, 3), then $a = \dots$

(a) zero

(b) 1

(c) - 1

(d) 3

(27) In any triangle XYZ, XY: YZ =

(a) sin X : sin Y

(b) $\sin Y : \sin Z$ (c) $\sin Z : \sin X$ (d) $\sin Z : \sin Y$

Second **Essay questions**

Answer the following questions:

Find algebraically in \mathbb{R} the solution set of : $\sqrt{x^2 - 4x + 4} = 10$

2 Find the value of: $\lim_{x \to 2} \frac{x^3 - 8}{x^2 - 5x + 6}$

Cairo Governorate



Nozha Educational Zone **Mathematics Inspection**

Multiple choice questions

Choose the correct answer from the given ones:



Interactive test 3

(1) The numerical value of the expression $\frac{\log 64}{\log 8}$ =

(a) 2

(c) 80

(d)72

(2) $\lim_{h \to 0} \frac{(x+h)^7 - x^7}{h} = \cdots$

(a) x^7

(b) $7 x^6$

(c) zero

(d) 1

- (3) In \triangle ABC: $a^2 + b^2 c^2 = \cdots$
 - (a) cos A

- (b) ab cos C
- (c) cos C
- (d) 2 ab cos C
- (4) If $5^{x} = 17$, then the value of x to the nearest two decimals =
 - (a) 1.34

- (b) 1.32
- (c) 1.76
- (d) 1.67

- (5) $\lim_{x \to \infty} \frac{2 X 1}{3 X + 1} = \dots$

- (b) $\frac{3}{2}$
- (c) ∞
- $(d) \infty$
- (6) The radius length of the circumcircle of Δ XYZ in which $\mathcal{X} = (20 \sin X) \text{ cm}$. equals cm.
 - (a) 5


- (b) 10
- (c) 20
- (d) 40

- (7) If $f(x) = 5^x$, then $f(-2) = \cdots$
 - (a) 2

- (c) $\frac{1}{25}$
- (d) $\frac{1}{5}$
- (8) The range of the function $f: f(X) = \frac{X-2}{2-X}$ equals
 - (a) IR

- (b) $\mathbb{R} \{2\}$
- (c) $\mathbb{R} \{-2\}$
- (9) All the following relations represent function y in terms of X except
 - (a) y = 3 X + 1
- (b) $y = x^2 4$ (c) $x = y^2 2$
- (d) $y = \sin x$
- (10) The value of $\log_5 49 \times \log_8 5 \times \log_9 8 \times \log_7 9 = \cdots$
 - (a) log 100
- (b) log 7
- (c) log 5
- (d) log 2

- (11) The rule of the function represented in the opposite figure is $f(x) = \cdots$
 - (a) $(x + 2)^2 + 3$
 - (b) $-(x-2)^2+3$
 - (c) $(x-1)^2 + 3$
 - (d) $(x+2)^2-3$



- (12) the logarithmic form that is equivalent to the exponential form: $2^7 = 128$ is
 - (a) $\log_2 128 = 7$

- (b) $\log_2 7 = 128$ (c) $\log_7 128 = 2$ (d) $\log_7 2 = 128$
- (13) The function $f: f(X) = a^X$ is increasing if
 - (a) a > 0

- (b) a > 1
- (c) a = 1
- (d) 0 < a < 1
- (14) The solution set in \mathbb{R} of the equation : |x-7| = 2 is
 - (a) $\{9,5\}$
- (b) $\{7,3\}$ (c) \emptyset
- (d) $\{3, -3\}$
- (15) In \triangle ABC, m (\angle C) = 61°, m (\angle B) = 71°, b = 91 cm., then $a \simeq \cdots \sim cm$.
 - (a) 71

- (b) 72
- (c)84
- (d) 92

(16)
$$\lim_{x \to 1} \frac{x^5 - 1}{x - 1} = \dots$$
(a) 5

(17) The curve of the even fun
(a) $y = x$

(18) If $\lim_{x \to 1} \frac{\sqrt{x} - 1}{x - 1}$

(b) 1

(c)4

(d) 20

(17) The curve of the even function is symmetric about the straight line

(b) vy

(c) xx

(d) y = -x

(18) If $\lim_{x \to 16} \frac{\sqrt{x-1}}{x-16} = \dots$

(a) zero

(b) $\frac{1}{2}$

(c) 1

(d) does not exist.

(19) If the perimeter of \triangle ABC = 33 cm., $\sin A + \sin C = \frac{2}{3}$, $\sin B = \frac{1}{4}$, then $AC = \cdots cm$.

(a) 6

(b)9

(c) 12

(d) 15

(20) If $\log 3 = X$, $\log 5 = y$, then $\log 15 = \dots$

(c) X + y

(d) X - y

(21) In \triangle ABC, $\frac{a}{a+b} = \frac{\sin A}{\cdots}$

(a) sin B

(b) sin C

 $(c) \sin A + \sin B$

(d) $\sin A + \sin C$

(22) The solution set of the equation : $\log_2 x \times \log_3 2 = 1$ in \mathbb{R} is

(b) $\{5\}$

(c) $\{4\}$

 $(d) \{3\}$

(23) $\lim_{x \to 0} \frac{x^7 - 1}{x + 1} = \dots$

(c) 1

(d) - 1

(24) The solution set of the following equation in \mathbb{R} : $\log_{\chi} 81 = 4$ equals

(c) $\{3, -3\}$ (d) $\{9\}$

(25) $\lim_{x \to a} \frac{ax}{3} = 12$, then $a = \dots$

 $(a) \pm 12$

(c) 3

(26) In \triangle ABC, $m(\angle A) : m(\angle B) : m(\angle C) = 3 : 5 : 4, then <math>c^2 : a^2 = \cdots$

 $(a) \sqrt{6:2}$

(b) 2:3

(c) 4:3

(27) $\lim_{x \to a} \frac{2x-4}{x-2} = \cdots$

(a) 1

(b) 2

(c) - 2

(d) zero

Second Essay questions

Answer the following questions :

If Graph the function $f: f(X) = X^2 + 1$, from the graph, deduce the range and it's monotony determine it's type whether it is even , odd or otherwise.

Find the value of: $\lim_{x \to 2} \frac{x^2 - x - 2}{x - 2}$

Giza Governorate



Awseem Educational directorate **Mathematics Inspection**

Multiple choice questions First



test (4)

Choose the correct answer from the given ones:

- (1) The domain of the function $f(x) = \sqrt{x-5}$ in \mathbb{R} is
 - (a) [5,∞[

- (b) $]5, \infty[$ (c) $]-\infty, 5]$ (d) $]-\infty, 5[$
- (2) XYZ is a triangle in which X = 4 cm., y = 8 cm., $m (\angle Z) = 75^{\circ}$, then $z = \cdots cm$.
 - (a) 6

- (b) 7
- (c) 8
- (d)9
- (3) The solution set of the equation : |X| + 5 = 0 in \mathbb{R} is
 - (a) $\{5\}$

- (b) $\{-5\}$ (c) $\{0\}$
- $(d) \emptyset$

2

-2 -3

-2 -1 O

(4) In the opposite figure:

$$\lim_{x \longrightarrow 2} f(x) = \cdots$$

- (a) 3
- (b) 2
- (c) 1
- (d) does not exist.

- (h) 49
- (d) zero

- (5) If $2^{X+1} = 5^{X+1}$, then $7^{X+1} = \dots$
 - (a) 1

- (6) The vertex of the function $f(x) = (x+1)^2 3$ is
 - (a) (1, -3)
- (b) (-1, -3) (c) (1, 3)
- (d)(-1,3)

- (7) If $\lim_{x \to \infty} \frac{a x^2 5 x}{3 x + 2 x^2} = 4$, then $a = \dots$
 - (a) 4

- (b) 2
- (c) 6
- (d) 8
- (8) The solution set of the equation : $\log_{\chi} 81 = 4$ in \mathbb{R} is
 - (a) $\{-3\}$

- (b) $\{3\}$
- (c) $\{3, -3\}$ (d) $\{9\}$
- (9) The radius length of the circle passing through the vertices of triangle XYZ in which y = 10 cm., $m (\angle Y) = 30^{\circ} \text{ equals} \cdots \text{cm.}$
 - (a) 40

- (b) 20
- (c) 10
- (d)5

- (10) $\lim_{x \to 1} \frac{x^7 1}{x 1} = \dots$
 - (a) 7

- (b) 1
- (c) 4
- (d)42

(11) The	axis of symmetry of	the function $f(X) = 1$	$\chi^2 + 3$ is the straight	line	
	x = 3	(b) $X = 0$		(d) y = 0	
(12) If lo	$g 2 = X, \log 5 = y,$		SA:		
(a) 2		(b) X – y		$(d)\frac{x}{y}$	
(13) If a	$\subseteq \mathbb{R}$ and $\lim_{x \to \infty} \frac{(a+3)}{2}$	$\frac{3) x^3 - 4 x^2 + 4}{x^2 + 5 x - 1} = -2$	2 , then a =	J	
(a) -	-7	(b) - 3	(c) zero	(d) 3	
(14) In tr	iangle ABC if 3 sin A	$a = 4 \sin B = 6 \sin C$	then $a:b:c=\cdots$		
	: 3:4			(d) 6:4:3	
(15) The	S.S. of the equation:	$\chi^{\frac{2}{3}} = 25$ in \mathbb{R} is			
(a) {	5}	(b) $\{-5,5\}$	(c) {125}	(d) {-125,125}	
(16) log ₅	$125 + \log 10 + \log_3 ($	25 + 2) =		Les 9	
(a) 3		(b) 5	(c) 7	(d) 9	
(17) In tri	angle ABC if $a = 3$ cr	m., $b = 5 \text{ cm}$.	$m (\angle C) = 100^{\circ}$	X-2-2	
	n the area of triangle				
(a) 5		(b) 6	(c) 7	(d) 8	
(18) $\lim_{x \to \infty} x$	$\frac{x^2 + 2x - 1}{3x^2 + 1} = \dots$	*******			
(a) $\frac{2}{3}$	_	(b) $\frac{3}{2}$	(c) $\frac{1}{2}$	(d) 2	
(<mark>19</mark>) If y =	= f(X) is the curve of	a real function, the	n its image by a trans	slation of magnitude	
2 units to the left is g $(x) = \cdots$					
1200 1000	(X+2)		(c) $f(x) + 2$	$ (\mathbf{d}) f(\mathbf{X}) - 2 $	
$(20) \lim_{x \to \infty}$	$\int_{3}^{1} \frac{x^2 - 6x + 9}{x - 3} = \dots$				
(a) ze	ero	(b) 3	(c) 6	(d) does not exist.	
(21) In tria	angle ABC , if m (∠	B) = 30° , c = $12\sqrt{3}$			
(a) 6°	$\sqrt{3}$	(b) 6	(c) 9	(d) 12	
22) The f	unction $f(X) = a^X$ is	increasing if	\$0.50		
(a) a :	> 0	(b) $a > 1$	(c) $a = 1$	(d) $0 < a < 1$	
23) Lim	$\frac{(3+h)^4-81}{h} = \cdots$				
(a) 4	•	(b) 81	(c) 108	(d) does not exist.	
24) If AB	24) If ABCD is a cyclic quadrilateral, then cos B + cos D =				
(a) ze		(b) 1	1	(d) - 1	

(25) If $\log_3 (2 X + 3) = 2$, then $X = \dots$

(a) 2

- (c) 4
- (d) 9

(26) If $\lim_{x \to \infty} (2x^{-3} + 3x^{-4} + a) = 3$, then $a = \dots$

(a) 6

- (c) 3
- (d) ∞

(27) The S.S. of the equation : $3^{X+1} + 3^X = 12$ in \mathbb{R} is

(a) $\{0\}$

- (b) $\{1\}$ (c) $\{1,0\}$
- $(d) \{2\}$

Essay questions Second

Answer the following questions:

Draw the graph of the function $f(x) = (x-1)^2 + 2$ and from the graph find range of the function and discuss its type for being even, odd or neither.

If $\lim_{x \to a} \frac{x^6 - a^6}{x^5 - a^5} = \frac{18}{5}$, then find the value of a

Giza Governorate



Aguzza Educational Zone

Multiple choice questions First



Choose the correct answer from the given ones:

- (1) The domain of the function $f(x) = \sqrt{x+2}$ is
 - (a) $[-2, \infty]$
- (b) $]-\infty, -2[$ (c) $\mathbb{R}-\{-2\}$
- (d) R
- (2) If $\log_a (X-3) + \log_a (X) = \log_a 4$, then $X = \cdots$
 - (a) 1

- (d) 1, 4
- (3) All the functions defined by the following rules are even except

- (b) $\chi^2 \sec \chi$ (c) $7 \chi^2 + 5$
- (4) The solution set of the equation : $\chi^{\frac{6}{5}} 64 = 0$ in \mathbb{R} is
 - (a) $\{-32, 32\}$
- (b) {32}
- (c) $\{-8, 8\}$
- (5) If $f(x) = 3^{x}$, then the solution set in \mathbb{R} of the equation : f(x-2) + f(x-1) = 36is
 - (a) {9}

- (b) $\{4\}$
- (c) $\{2\}$
- $(d) \{3\}$

(6) The curve g (X) = |X + 3| is the same as the curve f(X) = |X| by translation 3 units in the direction of

(a) OX

- (b) OX
- (c) OY
- (d) OY

- (7) The solution set of the inequality: $|3-2x| \le 1$ in \mathbb{R} is
 - (a) [1,2]

- (b)]1,2[(c) \mathbb{R}]1,2[(d) \mathbb{R} [1,2]
- (8) The range of the function $f: f(x) = x^2$ is
 - (a) [0,∞[
- (b)]0,∞
- (c) $]-\infty,0]$ (d) $]-\infty,0[$
- (9) The point of symmetry of the function $f: f(x) = \frac{2x-1}{x}$ is
 - (a)(1,1)
- (b) (2, 1)
- (c)(1,2)
- (d)(0,2)

- (10) If $5^{X-7} = 4^{7-X}$, then $X = \dots$
 - (a) $\frac{5}{4}$

- (d) zero
- (a) $\frac{5}{4}$ (b) 7 (c) $\frac{4}{5}$ (11) The numerical value of the expression $\frac{\log a^b}{\log a^c} = \frac{1}{\log a^c}$ (c) $\frac{a^b}{c}$

- (d) b c

- (12) If $\sqrt[3]{x^2} = 9$, then $x = \dots$
 - (a) 27

- $(c) \pm 3$
- $(d) \pm 27$

- (13) If $3^{2} = 5$, then $9^{x} = \dots$
 - (a) 10

- (c) 5
- (d) 5

- (14) If $\lim_{x \to 1} \frac{2x + a}{x + 1} = 5$, then $a = \dots$
 - (a) 2

- (b) 5
- (c) 8
- (d) 10

(15) In the opposite figure:

$$\lim_{x \longrightarrow 2} f(x) = \cdots$$

- (a) 2
- (b) 3
- (c) 1
- (d) does not exist.
- (16) $\lim_{x \to 1} \frac{2x-4}{x-2} = \dots$
 - (a) 1

- (c) 2
- (d) zero

-2-

- (17) If $\lim_{x \to \infty} \frac{10 x^m 2 x + 3}{7 x 2 x^3 + 1} = -5$, then m =

- (c) 1
- (d) 5

- (18) $\lim_{x \to 1} \frac{b}{x+1} = 5$, then $b = \dots$
 - (a) 5

- (b) 1
- (c) 1
- (d) 10

x'

- (19) $\lim_{x \to 1} \frac{2x+1}{\sqrt{x+3}-1} = \dots$

- (b) 3
- (c) 2
- (d) does not exist.

- $(20) \lim_{x \longrightarrow 0} \frac{\pi}{4} = \cdots$

- (b) 45
- (c) $\frac{\pi}{4}$
- (d) does not exist.

- (21) $\lim_{x \to \infty} (2)^{\frac{3}{x}} = \dots$
 - (a) zero

- (c) 1
- (d) 2
- (22) In $\triangle XYZ$, $3 \sin X = 4 \sin Y = 2 \sin Z$, then $X : y : z = \cdots$
 - (a) 2:3:6
- (b) 6:2:3
- (c) 4:3:6
- (23) In triangle ABC, $m (\angle A) = 45^{\circ}$, then length of the radius of its circumcircle = 6 cm. then $a = \cdots cm$.
 - (a) 13

- (b) $6\sqrt{2}$
- (c) 12
- $(d)\sqrt{2}$

- (24) In triangle ABC, $\frac{a}{a+b} = \frac{\sin A}{\dots}$
 - (a) sin B

- (b) sin C
- (c) sin A + sin B
- (d) $\sin A + \sin C$
- (25) In \triangle ABC, $\frac{b}{2 r} = \cdots$ where r is the length of the radius of the circumcircle of Δ ABC
 - (a) sin B

- (b) $\sin (A + B)$
- (c) $\sin A + \sin B$
- (d) sin A
- (26) In \triangle ABC, m (\angle A) = 112°, m (\angle B) = 33°, c = 19 cm. , then $b \simeq \dots$ to the nearest cm.
 - (a) 16

- (b) 17
- (c) 18
- (d) 20

(27) In the opposite figure:


The length of $AB \simeq \cdots$ to the nearest cm.

(a) 6

(b) 7

(c) 8

(d)9



Essay questions Second

Answer the following questions:

- 1 Find: $\lim_{x \to -3} \frac{x^6 729}{x^3 + 27}$
- 2 Draw the curve of the function : $f(x) = 3 (x 1)^2$, from the curve : determine the domain and the range, discuss its monotonicity, determine the type of the function (even , odd or otherwise)

Alexandria Governorate



Math Inspection

First Multiple choice questions



test 6

Choose the correct answer from the given ones:

(1) The domain of the function $f: f(x) = \sqrt{x-3}$ is

(a) IR

- (b) $\mathbb{R} \{3\}$ (c) $[3, \infty[$
- (d) $]-\infty$, 3
- (2) The range of the function $f: f(x) = \begin{cases} 0 & \text{is } x \le 0 \\ 1 & \text{is } x > 0 \end{cases}$
 - (a) $\{1\}$

- (b) {0}
- (c) IR
- (d) $\{0,1\}$

- (3) The type the function $f: f(x) = \frac{\sin x}{x}$ is
 - (a) even.

(b) odd.

(c) neither even nor odd.

- (d) linear.
- (4) The point of symmetry of the curve of the function $f: f(x) = \frac{1}{x-3} + 4$ is
 - (a) (3, -4)
- (b) (-3, -4)
- (c) (3,4)
- (d) (-3,4)
- (5) The solution set of the equation: $|2 \times -4| = |\times +1|$ is
 - (a) $\{1,5\}$

- (b) $\{5, -1\}$ (c) $\{1, -5\}$ (d) $\{-5, -1\}$
- (6) If $3^{x-5} = 9$, then $x = \dots$
 - (a) 7

- (b) 3
- (c) 2

- $(7)^4 \sqrt{x^4 y^8} = \cdots$
 - (a) χv^2

- (b) $|X|y^2$ (c) $\pm Xy^2$
- (d) $\chi |v^2|$

- (8) If $f(x) = (5)^{-x}$, then $\frac{f(x-1)}{f(x+1)} = \dots$

- (b) $\frac{1}{5}$
- (d) $\frac{1}{25}$
- (9) An amount of 5000 pounds is deposited in a bank gives a yearly compound interest 5 % for 7 years ≈ ····· pounds.
 - (a) 6750

- (b) 7035.5
- (c) 5350
- (d) 8500
- (10) The solution set of the equation $\log_{\mathcal{X}} (3 \ \mathcal{X} 2) = 2$ in \mathbb{R} is
 - (a) $\{1,2\}$
- (b) {1}
- $(c) \{2\}$
- (11) The curve of the function $f: f(X) = \log_2 X$ is passing through the point (8,)
 - (a) 2

- (b) 3
- (c) log₂ 3
- (d) 256

(12) If $X = 5 + 2\sqrt{6}$, then $\log \left(X + \frac{1}{X}\right) = \dots$

(a) 1

(b) $5 - 2\sqrt{6}$

(c) 10

(d) $5 + 2\sqrt{6}$

(13) If $3^{x} = 5$, then $x = \dots$

(a) 3

(b) log₃ 5

 $(c) \log_5 3$

(d) $\frac{5}{3}$

(14) $\lim_{x \to 2} \frac{x^2 + x - 6}{x^2 - 4} = \dots$ (a) $\frac{4}{5}$

(c) $\frac{2}{5}$

 $(d) - \frac{2}{5}$

(15) $\lim_{x \to 0} \frac{\sqrt{x+1}-1}{x} = \cdots$

(a) zero

(c) $\frac{1}{2}$

(d) not exist.

(16) If $\lim_{x \to 2} \frac{x^2 - 4a}{x - 2}$ exist, then $a = \dots$

(c) 2

(d)4

(17) $\lim_{x \to -2} \frac{x^7 + 128}{x^4 - 16} = \dots$

(c) - 14

(d) 14

(18) If $\lim_{x \to -k} \frac{x^5 - k^5}{x - k} = 80$, then $k = \dots$

 $(c) \pm 2$

(d) 16

(a) 2 (b) -2 (19) The $\lim_{x \to \infty} \frac{x^7 - 2x^3}{2x^4 - 3x^2 - 1} = \dots$

(c) ∞

(d) $\frac{1}{2}$

(20) $\lim_{x \to \infty} \frac{\sqrt{8+9 x^2}}{x} = \cdots$

(a) $2\sqrt{2}$

(b) 3

(c) $-2\sqrt{2}$

(d) - 3

(21) $\lim_{x \to \pi} \frac{\cos 2 x}{x} = \cdots$

(b) 1

(c) $\frac{1}{\pi}$

(22) DEF is a triangle in which m (\angle D) = 80° and m (\angle E) = 60°, if f = 12 cm. • then d = cm.

(a) $\frac{12 \sin 80^{\circ}}{\sin 40^{\circ}}$

(b) $\frac{12 \sin 80^{\circ}}{\sin 60^{\circ}}$ (c) $\frac{12 \sin 40^{\circ}}{\sin 80^{\circ}}$ (d) $\frac{12 \cos 80^{\circ}}{\cos 40^{\circ}}$

(23) ABC is a triangle in which $\frac{\sin A}{3} = \frac{\sin B}{5} = \frac{\sin C}{4}$, then a: b: c =

(a) 6:5:8

(b) 8:5:6

(c) 7:2:4

(d) 3:5:4

(24) In \triangle LMN, $\ell = 5$ cm., m = 7 cm., $m (\angle N) = 60^{\circ}$, then $n = \cdots$ cm. (to the nearest tenth)

(a) 6.2

(b) 5

(c) 4.3

(d) 3.5

- (25) In \triangle XYZ, $y^2 + z^2 x^2 = 2$ y z ×
 - (a) cos X

- (b) sin Z
- (c) cos Z
- (d) sin X
- (26) The number of possible solutions of \triangle XYZ in which X = 5 cm. y = 6 cm. , m ($\angle X$) = 70° equals
 - (a) zero

- (c) 1
- (d) 3
- (27) If ABC is a triangle in which $\frac{2}{\sin A} = \frac{3}{\sin B} = \frac{4}{\sin C}$, then the measure of the smallest angle in the triangle ≈ ······
 - (a) 57° 28

- (b) 41° 12 (c) 28° 57
- (d) 36° 52

Second **Essay questions**

Answer the following questions:

- Find the following limit (show your steps): $\lim_{x \to -3} \frac{x^3 10 x 3}{x^2 + 2 x 3}$
- **2** Find algebraically in \mathbb{R} the solution set of the following inequality: |x-5| > 3
 - El-Kalyoubia Governorate



Math Inspection

Multiple choice questions First



test (7)

Choose the correct answer from the given ones:

- (1) The solution set of the equation |X| + 4 = 0 in \mathbb{R} is
 - (a) $\{-4\}$

- (b) {2}
- $(d) \emptyset$
- (2) The domain of the function $f(x) = \frac{5}{\sqrt{x-1}}$ is
 - (a) R+

- (b) $\mathbb{R} \{1\}$
- (c) $\mathbb{R} \{0\}$ (d) $[1, \infty[$
- (3) The point of symmetry of the function $f(x) = \frac{-1}{x-3} + 4$ is
 - (a) (3, 4)

- (b) (-3,4)
- (c) (3, -4)
- (d) (-3, -4)
- (4) The solution set of the inequality: $\sqrt{\chi^2} 9 < 0$ in \mathbb{R} is
 - (a) [-3,3]
- (b)]-9,9[(c) $\{3\}$
- (d) $\{9\}$
- (5) The range of the function f(x) = |x-2| + 3 is
 - (a) $]-\infty$, 2
- (b) $[2, \infty[$ (c) $[3, \infty[$
- $(d) \left[-2, \infty \right[$

(6) If
$$\left(\frac{2}{5}\right)^{X-3} = \frac{5\sqrt{5}}{2\sqrt{2}}$$
, then $X = \dots$
(a) $\frac{3}{2}$ (b) $\frac{9}{2}$

(a)
$$\frac{3}{2}$$

(b)
$$\frac{9}{2}$$

(7)
$$\log_{\sqrt{2}} 4 \times \log_{\sqrt{2}} 2 = \log_{\sqrt{2}} \dots$$

(8) If
$$\log 3 = X$$
 and $\log 4 = y$, then $\log 12 = \dots$

(a)
$$X + y$$

(c)
$$\chi - y$$

(d)
$$\log x + \log y$$

(9) If
$$f(x) = b^x$$
 is passes through the point (2, 4), then $b = \dots$

(b)
$$\frac{1}{2}$$

(d)
$$\frac{1}{4}$$

(10) If
$$\log_2 x = 3$$
, then $\log_8 x = \cdots$

(a)
$$\frac{3}{2}$$

(b)
$$\frac{2}{3}$$

(11) If
$$\sqrt[3]{x^2} = 4$$
, then $x = \dots$

$$(c) \pm 8$$

$$(d) \pm 16$$

(12)
$$\log_2 12 + \log_2 \frac{2}{3} = \cdots$$

(13) If
$$4^{X-2} = 3^{2X-4}$$
, then $X = \dots$

(14)
$$\lim_{x \to 2} 5 = \cdots$$

(15)
$$\lim_{X \to -2} \frac{1}{X} = \dots$$

$$(b) - 1$$

(c)
$$\frac{1}{2}$$

(d)
$$\frac{-1}{2}$$

(16)
$$\lim_{x \to 1} \frac{x^5 - 1}{x^3 - 1} = \cdots$$

(a)
$$\frac{5}{3}$$

(b)
$$\frac{-5}{3}$$

$$(d) - 15$$

(17)
$$\lim_{X \to \infty} \frac{3-X}{X} = \cdots$$

$$(a) - 1$$

(18)
$$\lim_{x \to 1} \frac{x^2 - 1}{x - 1} = \dots$$

$$(d)$$
 3

(19)
$$\lim_{x \to \infty} \frac{(a-2) x^4 + b x^3 - 5}{3 x^3 + 7} = \frac{1}{3}$$
, then $a + b = \dots$

$$(d)$$
 5

(20)
$$\lim_{x \to -3} \frac{x^3 + 27}{x^2 - 9} = \dots$$

(a) $\frac{3}{2}$ (b) $\frac{-9}{2}$
(21) $\lim_{x \to \frac{\pi}{2}} \frac{\sin x}{x} = \dots$

(a)
$$\frac{3}{2}$$

(b)
$$\frac{-9}{2}$$

(c)
$$\frac{2}{3}$$

(d)
$$\frac{1}{2}$$

$$(21) \lim_{X \longrightarrow \frac{\pi}{2}} \frac{\sin X}{X} = \dots$$

(b)
$$\frac{\pi}{2}$$

$$(c)\frac{2}{\pi}$$

(d)
$$\frac{1}{90^{\circ}}$$

(a) 1 (b)
$$\frac{\pi}{2}$$
 (c) $\frac{2}{\pi}$ (22) In \triangle ABC, if $\frac{a^2 + b^2 - c^2}{2 a b} = 0$, then

(a) m (
$$\angle A$$
) = 60°

(b) m (
$$\angle$$
 B) = 90°

(c) m (
$$\angle$$
 C) = 120°

(d) m (
$$\angle$$
 A) + m (\angle B) = 90°

(23) In
$$\triangle$$
 XYZ, if $X = y = 8$ cm. and the perimeter of \triangle XYZ = 26 cm., then m (\angle Z) \simeq

(24) In
$$\triangle$$
 ABC, if $a = 3$ cm., $b = 4$ cm., $c = 6$ cm., then $\cos C = \cdots$

(a)
$$\frac{-11}{24}$$

(b)
$$\frac{-11}{12}$$
 (c) $\frac{11}{24}$

(c)
$$\frac{11}{24}$$

(d)
$$\frac{11}{12}$$

(25) In
$$\triangle$$
 ABC, if $a = 6$ cm., $m (\angle B) = 2$ m $(\angle A) = 80^{\circ}$, then $c = \cdots$

$$\frac{6\sin 40^{\circ}}{\sin 60^{\circ}}$$

(b)
$$\frac{\sin 60}{6 \sin 40^{\circ}}$$
 (c) $\frac{\sin 40^{\circ}}{6 \sin 60^{\circ}}$ (d) $\frac{6 \sin 60^{\circ}}{\sin 40^{\circ}}$

$$\frac{\sin 40^{\circ}}{6 \sin 60^{\circ}}$$

$$\frac{\text{(d)}}{\sin 40^{\circ}} \frac{6 \sin 60^{\circ}}{\sin 40^{\circ}}$$

(c)
$$2\sqrt{3}$$

(d)
$$4\sqrt{3}$$

(27) In
$$\triangle$$
 ABC, $\frac{1}{\sin B + \sin C} = \frac{a}{\sin A}$

(b)
$$b + c$$

(c) area of
$$\triangle$$
 ABC

(d) The perimeter of
$$\triangle$$
 ABC

Second Essay questions

Answer the following questions :

Use the curve of $f(X) = \frac{1}{X}$ to represent the curve of g(X) = 2 + f(X - 1), from the graph determine the domain and the range.

2 Find:

$$(1) \lim_{x \to 5} \frac{x-5}{\sqrt{x-1}-2}$$

$$(2) \lim_{x \to \infty} \frac{x+2}{\sqrt{9x^2+25}}$$

El-Monofia Governorate



Menouf Educational Administration Mathematic Inspection

Multiple choice questions



(1) The domain of the function $f(x) = \sqrt{x-4}$ is

- (a) [4,∞
- (b)]-∞,4[(c)]4,∞[
- (d) $-\infty$, 4
- (2) The function which is even from the functions defined by the following rules $f(X) = \cdots$
 - (a) $X \cos X$
- (b) $X \sin X$ (c) $X^3 + 1$
- (d) $\tan x$

- (3) The range of function f(X) = -|X| is
 - (a) R

- (b) $]0, \infty[$ (c) $]-\infty, 0[$ (d) $]-\infty, 0]$
- (4) The symmetric point of the function $f: f(x) = x^3 1$ is
 - (a)(0,1)
- (b) (0,-1) (c) (-1,1) (d) (0,0)
- (5) The function $f(x) = (x-2)^2 + 3$ is increasing on the interval
 - (a) IR

- (b) $]2, \infty[$ (c) [-2, 2] (d) $]-\infty, 2[$
- (6) If $3^{x-5} = 9$, then $x = \dots$
 - (a) 2

- (b) 3
- (c) 5
- (d)7
- (7) The solution set of the equation $3^{x+1} + 3^x = 12$ in \mathbb{R} is
 - (a) $\{0\}$

- (b) {1}
- (c) $\{3\}$
- (d) $\{0,1\}$
- (8) The exponential function of base a is increasing if
 - (a) a > 0

- (b) a > 1
- (c) 0 < a < 1
- (d) a = 1
- (9) An amount of 5000 pounds is deposited in a bank gives a yearly compound interest 5 % for 7 years ≈ ····· pounds.
 - (a) 5350

- (b) 6750
- (c) 7035.5
- (d) 8500

- (10) If $\log_3 x = 2$, then $x = \cdots$
 - (a) 3

- (b) 5
- (c) 8
- (d)9

- (11) $\log_2 5 \times \log_5 2 = \cdots$
 - (a) 1

- (b) 10
- (c) log 10
- (d) log 7

- (12) If $\log x + \log 5 = 2$, then $x = \dots$
 - (a) 3

- (b) 8
- (c) 17
- (d) 20

(13) The	e solution set of the equ	nation $\log_{\chi} (64 \ X) =$	4 in R is	
	{2}	20 2 200	(c) {0,4}	(d) {6}
(14) Li	$\lim_{x \to 4} \left(3 \ X - \sqrt{X} \right) = \dots$	ronne		11200 114 12
(a)		(b) 10	(c) 14	(d) 16
(15) Li	$\underset{\rightarrow}{\text{m}} \frac{\chi^3 - 8}{\chi^2 - 4} = \dots$			
(a)	2	(b) 3	(c) 4	(d) 6
(16) Li	$\underset{\rightarrow}{\text{m}} \frac{\sqrt{x+1}-2}{x-3} = \dots$			
(a)	$\frac{1}{4}$	(b) 4	(c) $\frac{1}{6}$	(d) 6
(17) Li				
(a)	zero	(b) $\frac{5}{3}$	(c) 4	(d) $6\frac{2}{3}$
(18) Li				J J
(a)	zero	(b) 2	(c) 4	(d) 8
(19) Li	$\underset{\bullet}{\text{m}} \frac{(X+1)^9 - 1}{X} = \dots$			
(a)	zero	(b) 1	(c) 9	(d) 10
(20) Li	$\underset{+\infty}{\text{m}}$ (3 χ^{-5} + 4 χ^{-2} + 5	5) =		
	zero	(b) 5	(c) 12	(d) ∞
(21) If x	$\lim_{x \to \infty} \frac{a x + 6}{2 x - 7} = 4, \text{ then}$	a =		
(a) 2	2	(b) 4	(c) 6	(d) 8
	ABC if m (\angle A) = 60° umcircle = 5 cm. then t			
(a) 9	9	(b) 12	(c) 31	(d) 62
(23) If A	BCD is a cyclic quadril	lateral, then cos A+	cos C =	
(a) 1	1	(b) zero	(c) $\frac{1}{2}$	(d) - 1
(24) In Δ	XYZ, then $2 \times z \times \cdots$	$\cdots\cdots\cdots = x^2 + z^2 - z^2$	y^2	
(a) (cos X	(b) cos Y	(c) cos Z	(d) sin Y
	ABC if a = 4 cm. , b = BC =	$7 \text{ cm.}, \text{m} (\angle C) = 1$	20°, then the area of	of
(a) 7		(b) 7√3	(c) 14	(d) $14\sqrt{3}$

		77207	1000	
Final	exam	ning	tion	C
11101	CAGII	11110	шил	

(26) In \triangle ABC, m (\angle A) = 30°, a = 7 cm., then the length of diameter of its circumcircle is

(a) 7

- (b) $7\sqrt{2}$
- (c) 14
- (d) $14\sqrt{2}$

(27) In \triangle ABC if sin A: sin B: sin C = 3:4:2, then m (\angle C) \approx nearest degree.

(a) 29

- (b) 57
- (c) 82
- (d) 89

Second Essay questions

Answer the following questions:

- Graph the function $f: f(x) = (x-2)^2$ determine its type whether it is even, odd or otherwise and deduce the range.
- If $\lim_{x \to a} \frac{x^8 a^8}{x^5 a^5} = 25$ find the value of a
 - 9 El-Dakahlia Governorate



Maths Supervision

First Multiple choice questions



Choose the correct answer from the given ones:

Interactive test 9

- (1) In any triangle XYZ, $z: x = \dots$
 - (a) sin X: sin Y
- (b) sin Y : sin Z
- (c) sin Z : sin X
- (d) sin Z: sin Y
- (2) In \triangle ABC, if $\frac{\sin A}{4} = \frac{\sin B}{9} = \frac{\sin C}{7}$, then the greatest angle in measure is
 - (a) ∠ A

- (b) ∠ B
- (c) \(\text{C}
- (d) right.
- - (a) a right-angled triangle.

(b) an isosceles triangle.

(c) an equilateral triangle.

- (d) a scalene triangle.
- (4) The radius length of the circumcircle of the triangle ABC in which m (\angle A) = 30°, and a = 10 cm. is cm.
 - (a) 5

- (b) 10
- (c) 20
- (d) 40
- - (a) 5

- (b) 10
- (c) 15
- (d) 20

(6)	In \triangle ABC, 6 a = 4 b = 3 d	then the measure	of the smallest angle	in triangle =
	(a) 57° 28	(b) 41° 12		(d) 36° 52
(7)	In all the following relation			
		(b) $y = X^2 - 4$		(d) $y = \sin x$
(8)	The domain of the functio			
	(a) R		(c) [3,∞[(d) $]-\infty,3]$
(9)	The type of function $f: f$	$(X) = \frac{\sin X}{x}$ is	8	
	(a) even.		(b) odd.	8
	(c) linear.		(d) neither even nor	odd.
(10)	The function $f: f(X) = $	$ \begin{array}{cccc} 2 & , & \chi > 0 \\ -2 & , & \chi < 0 \end{array} $ is syr	nmetric about the po	int
	(a) (2,0)	(b) $(-2,0)$	(c) (0 , 0)	(d) $(2, -2)$
(11)	The point of symmetry of	the curve of the func	tion $f: f(x) = \frac{1}{x-3}$	+ 4 is
	(a) (3,4)	(b) $(3, -4)$	(c)(-3,4)	(d) $(-3, -4)$
(12)	$\lim_{x \longrightarrow 3} 15 = \cdots$			
	(a) 4	(b) 45	(c) 15	(d) 18
(13)	$\lim_{x \longrightarrow 2} \frac{x^2 - 4}{x - 2} = \dots$			
	(a) undefined.	(b) 8	(c) - 4	(d) 4
(14)	$\lim_{x \to 0} \frac{\sqrt{x+1} - 1}{x} = \dots$;se:		
	(a) zero	(b) $\sqrt{2}$	(c) $\frac{1}{2}$	(d) has no existence.
(15)	If $\lim_{x \to 2} \frac{a}{x+1} = 4$, then a	=		
	(a) 3	(b) 4	(c) 12	(d) $\frac{2}{3}$
(16)	$\lim_{x \to 2} \frac{x^5 - 32}{x^3 - 8} = \dots$	超		
		(b) zero	(c) $6\frac{2}{3}$	(d) $\frac{5}{3}$
(17)	$\lim_{x \to 0} \frac{(x+1)^9 - 1}{x} = \dots$		3	3
	(a) 9	(b) 1	(c) zero	(d) 10
(18)	$\lim_{x \to \infty} \left(\frac{3}{x^2} - 2 \right) = \dots$			
	(a) 3	(b) 2	(c) - 3	(d) - 2
(19)	$\lim_{x \to \infty} \frac{2x^2 + 1}{x^2 + 1} = \dots$	10		
	(a) zero	(b) 2	(c) ∞	(d) doesn't exist

(20) $a^m \times a^m = \cdots$

(a) a^{m²}

(c) 2 a^m

(d) m a^2

(21) If $3^{x-5} = 9$, then $x = \dots$

(a) - 7

(b) - 3

(d)7

(22) Which of the following is not equal to $\sqrt[5]{x^4} = \cdots$ (a) $(\sqrt[5]{x})^4$ (b) $\sqrt[4]{x^5}$ (c) $x^{\frac{4}{5}}$

(a) $(\sqrt[5]{x})^4$

(b) $\sqrt[4]{x^5}$

(d) $(x^{\frac{1}{5}})^4$

(23) If the curve of the function $f: f(X) = a^X$ passing through (1,3), then $a = \cdots$

(b) 1

(c) - 1

(24) $\operatorname{Log}_{a} X = y$ is equivalent to

(a) $\log_a y = X$

(b) $a^y = X$

(c) $a^{\chi} = y$

(d) y = a X

(25) The solution set of the equation $\log_{\chi}(\chi + 6) = 2$ in \mathbb{R} is

(a) $\{3\}$

(b) $\{3, -2\}$ (c) $\{3, 1\}$ (d) $\{6, 1\}$

(26) The curve of the function $f: f(X) = \log_2 X$ passing through the point (8,)

(a) 2

(c) log₂ 3

(d) 256

 $\frac{1}{\log_2 14} + \frac{1}{\log_2 14} = \dots$

(b) 2

(c) 7

(d) 14

Second Essay questions

Answer the following questions:

11 Draw the curve of the function f where $f(X) = X^2$, $X \subseteq \mathbb{R}$, from graph determine the range and the type of the function (even, odd, or neither even nor odd)

2 Find: $\lim_{x \to 2} \frac{x^2 - 4}{x - 2}$

Damietta Governorate



Maths Inspection

First Multiple choice questions



Choose the correct answer from the given ones: (1) The domain of the function $f: f(x) = \sqrt{x-5}$ is

(a) [5,∞[

(b) $]-\infty, 5]$ (c) $]5, \infty[$

 $(d) \begin{bmatrix} -5, \infty \end{bmatrix}$

(2) Which of the functions that are defined by the following rules represents an exponential decay function?


(a) $f(x) = 2^x$

(b) $f(x) = \left(\frac{1}{3}\right)^{-x}$ (c) $f(x) = 3^x$ (d) $f(x) = \left(\frac{2}{3}\right)^x$

(3) In the opposite figure:

$$\lim_{x \to 2} f(x) = \dots$$

- (a) 3
- (b) 2
- (c) 1
- (d) does not exist.



- (4) A circle with diameter of length 20 cm., passes through the vertices of Δ ABC which is an acute-angled triangle in which BC = 10 cm., then m (\angle A) =°
 - (a) 30


- (b)60
- (c)45
- (d) 150
- (5) The even function from the functions that are defined by the following rules is
 - (a) $f(x) = x^3$

- (b) g(X) = 3 X (c) $h(X) = \frac{1}{x}$ (d) $n(X) = x \sin x$
- (6) If $2^{x-1} = 7$, then $x = \dots$

- (d) 3.6
- (7) If $\lim_{x \to 3} \frac{x^2 2x + k}{x^2 9} = m$, where $m \in \mathbb{R}$, then $k \times m = \dots$

- (d) 1
- (8) In \triangle ABC, $\frac{2 \text{ b}}{\sin B} = \dots \text{r}$ (where r is the radius of its circumcircle)
 - (a) 1

- (b) 2
- (c) 4
- (d) 8



- (10) If $\lim_{x \to 3} \frac{x^2 2x a}{x 3} = 4$, then $a = \dots$

- (b) 3
- (c) 1
- (d) 3
- (11) In \triangle ABC, a = 9 cm., b = 15 cm., m (\angle C) = 106° , then its perimeter \simeq cm.
 - (a) 44

- (b) 42
- (c) 34
- (d) 28

(12) The range of the function $f: f(X) = X $ is				
(a) R + 4/5	(b) ℝ [−]	(c) R	$(d) [0, \infty[$
$(13)_{x}$	a) \mathbb{R}^+ Lim $\frac{\sqrt[4]{x^5 - 32}}{x - 16} = \dots$			
(a) 5	(b) $\frac{5}{2}$	(c) $\frac{5}{4}$	(d) $\frac{5}{8}$
(14) T	The solution set of the equa	ation: X + 2 = 0 in	R is	
(a) $\{-3\}$	(b) {3}	(c) $\{-3,3\}$	(d) Ø
(15) le	$\log_b a \times \log_c b \times \log_d c \times \log_d c$	$og_a d = \cdots$		
(a) zero	(b) 1	(c) abcd	(d) ad
(16) _x	$\lim_{x \to \infty} (5 + 3^{\frac{1}{X}}) = \dots$	\$		
	(a) 8		(c) 5	(d) 6
(17) x	$\lim_{x \to 0} \frac{1 - \cos \theta}{3 - x} = \dots$			
	(a) $\frac{1}{3}$	(b) $\frac{1}{2}$	(c) 1	(d) zero
(18) I	In \triangle ABC, if $2 \sin A = 3 \text{ s}$	in $B = 4 \sin C$, then	a:b:c=	
((a) 4:6:3	(b) 6:3:4	(c) 3:6:4	(d) 6:4:3
(19)	The solution set of the equ	ation: $3^{x} + 3^{2+x} = 3^{2+x}$	90 in R is	
	(a) {1}		(c) {3}	(d) $\{-3\}$
(20) I	If $\log 3 = x$, $\log 4 = y$, th	nen log 12 =		
((a) $X + y$	(b) X y	(c) $X - y$	(d) $\log x + \log y$
(21) I	If ABC is a triangle in whi	ch a = 4 cm. $b = 4^{\circ}$	$\sqrt{3}$ cm., c = 8 cm.,	then cosine of the
	smallest angle equals	114		8
	(a) $\frac{1}{2}$	(b) $\frac{\sqrt{3}}{2}$	(c) 1	(d) zero
(22)	If $\lim_{x \to \infty} \frac{4 a x^n - 4 x + 5}{3 - 9 x + 8 x^2}$	$= 3$, then $a + n = \cdots$		
	(a) 8	(b) - 8	(c) 9	(d) 4
(23)	The straight line $y = 9$ cuts	s the curve of the fun	$ction f: f(X) = 3^X a$	at the point
	(a) (0,9)	(b) $(-2,9)$	(c) (2,9)	(d) (1,9)
(24)	$\lim_{X \to 3} \frac{(X-2)^7 - 1}{X-3} = \dots$			
9	(a) 7	(b) 14	(c) 2	(d) - 2
(25)	The solution set in \mathbb{R} of : \mathbb{V}	$\sqrt{x^2 - 6x + 9} < 5$ is		
9	(a)]-5,5[(b) $]-2,8[$	(c)]-8,2[(d) $]-2,5[$
				(d)

(26) In \triangle ABC, if m (\angle B) = 60°, m (\angle C) = 30°, c = 4 cm., then b = cm.

(a) 4

- (b) 8
- (c) $2\sqrt{3}$
- (d) $4\sqrt{3}$

(27) If $\log_3 (2 X + 3) = 2$, then $X = \dots$

(a) 3

- (b) 2
- (c) 9
- (d)4

Second Essay questions

Answer the following questions :

- 1 Draw the curve of the function $f: f(X) = \frac{1}{X-2} + 1$, then from the graph:
 - (1) Discuss the monotonicity of f
 - (2) Determine whether f is even, odd or otherwise.

2 Find: $\lim_{x \to 5} \frac{\sqrt{x-1}-2}{x-5}$

El-Beheira Governorate



Maths Inspection

First Multiple choice questions

Choose the correct answer from the given ones:

- (1) The range of the function f(x) = x 2 is
 - (a) R+

- (b) R -
- (c) $\mathbb{R} \{2\}$
- (2) The point of symmetry of the curve of the function $f(x) = x^3 + 1$ is
 - (a)(1,0)

- (b) (-1,0)
- (c)(0,1)
- (d)(0,-1)

- (3) If f(x) = 2, then $f(3) = \cdots$
 - (a) 2

- (b) 4
- (c) 0
- (d) 1
- (4) The solution set of the inequality: $|x| \le 2$ in \mathbb{R} is
 - (a) $]-\infty,2]$
- (b)]-2,2] (c) [-2,2]
- (d)]-2,2[
- (5) The domain of the function f(x) = x 4 is
 - (a) 4,∞
- (b) $]4, \infty[$ (c) $[-\infty, 3[$
- (d) R
- (6) The solution set of the equation: $\chi^{\frac{3}{2}} = 8$ in \mathbb{R} is
 - (a) {4}

- (b) $\{4, -4\}$ (c) $\{8\}$
- (d) $\{-8, 8\}$

- (7) If $3^{x+1} = 5^{x+1}$, then $7^{x+1} = \dots$
 - (a) 0

- (b) 1
- (c) 2
- (d)3

(8) If $f(X) = 3^X$, then $f(-1) = \cdots$

(a) - 1

(b) 1

(c) $\frac{1}{3}$

 $(d) \frac{-1}{3}$

(9) If $\log 3 = a$, $\log 5 = b$, then $\log 15 = \cdots$

(a) ab

(c) a + b

(d) a - b

(10) If $\log_2 X = 3$, then $X = \cdots$

(a) 2

(b) 3

(c) 8

(d) 9

(11) $\log_2 3 \times \log_3 4 \times \log_4 5 \times \log_5 8 = \dots$

(a) 1

(c) 3

(d)4

(12) The solution set of the equation : $\log_2 X \times \log_3 2 = 4$ is

(a) $\{81\}$

(b) $\{4\}$

(c) $\{3\}$

 $(d) \{2\}$

(13) If $5^{x} = 17$, then $x = \dots$ (to the nearest hundredth)

(a) 1.34

(c) 1.76

(d) 1.67

(14) $\lim_{x \to 0} \frac{x^5 - 1}{x + 1} = \dots$

(c) - 2

(d) 5

(15) $\lim_{x \to \infty} (3 + 5 x^2 + 3 x)$

(a) not exist.

(b) 5

(c) ∞

(d) 11

(16) $\lim_{x \to 0} (3) = \cdots$

(a) 0

(b) 1

(c)2

(d) 3

(17) $\lim_{x \to \infty} \frac{3 x^2 + 4}{x^2 + 5} = \cdots$

(a) 3

(b) 4

(c)5

(d) 1

(18) If $\lim_{x \to 2} \frac{a}{x+1} = 3$, then $a = \dots$

(a) 6

(b) 8

(c)9

(d) 12

(19) $\lim_{x \to 2} \frac{x^5 - 32}{x^2 - 4} = \dots$

(b) 40

(c)60

(d) 80

(20) $\lim_{x \to 2} \frac{x^2 - 4}{x - 2} = \dots$

(c)0

(d) 8

(21) $\lim_{x \to 0} \frac{\sqrt{x+4}-2}{x} = \dots$

(a) 0

(b) 1

(c) $\frac{1}{4}$

(d)4

- (22) In \triangle ABC if m (\angle A) = 30° and a = 6 cm., then $\frac{b}{\sin B}$ =
 - (a) 3

- (b) 6
- (c) 8
- (d) 12
- (23) In \triangle ABC if $2 \sin A = 3 \sin B = 4 \sin C$, then $a : b : c = \dots$
 - (a) 2:3:4
- (b) 4:3:2
- (c) 3:4:6
- (d) 6:4:3
- (24) The length of the diameter of the circumcircle of the triangle ABC in which b = 12 cm. and m (\angle B) = 90° iscm.
 - (a) 6

- (b) 40
- (c) 20
- (d) 12
- (25) In \triangle ABC, a: b: c = 3:7:5, then the measure of the greatest angle in the triangle ABC is°
 - (a) 60

- (b) 30
- (c) 90
- (d) 120
- (26) In \triangle ABC : a = 4 cm. , b = 7 cm. and m (\angle C) = 120° , then the area of the triangle =
 - (a) 14

- (b) $7\sqrt{3}$
- (c) 28
- (d) $7\sqrt{2}$
- (27) In \triangle ABC: a = 10 cm., b = 10 cm. and m (\angle C) = 120°, then $c = \cdots cm$.
 - (a) 12

- (b) 14
- (c) $10\sqrt{3}$
- (d) 7

Second **Essay questions**

Answer the following questions :

- **1** Find the solution set in \mathbb{R} of the equation : $|2 \times -3| = 1$
- **2** Find: $\lim_{x \to -3} \frac{(x+5)^4 16}{x+3}$

El-Menia Governorate



Mattay Educational Directorate

Multiple choice questions First

Choose the correct answer from the given ones:

- (1) The domain of the function f: f(x) = 7 is
 - (a) {7}

- (b) R
- (c) $\mathbb{R} \{7\}$ (d) $\mathbb{R} \{0\}$

- $(2) f(X) = X + X^3 = \cdots$
 - (a) even.

(b) odd.

(c) neither even nor odd.

(d) anything else.

(3) Symmetric point $f(X) = X^3 - 1$ is

(a) (0, 0)

(b) (1,0) (c) (0,1) (d) (0,-1)

(4) Solve the equation : |x + 2| + 1 = 0 is

(a) R

(b) $\{3\}$ (c) $\{-1\}$

 $(d) \emptyset$

(5) Solve the equality: |X| < 2 is

(a) Ø

(b) $\mathbb{R} - [-2, 2]$ (c)]-2, 2[(d) [-2, 2]

(6) $2^{X+1} = 8$, then $X = \dots$

(a) 8

(b) 2

(c) 3

(d) 4

(7) $\lim_{x \to 1} (7x + 3) = \cdots$

(a) 7

(b) 3

(c) 10

(d) 1

(8) If the curve of function $f: f(x) = 5^x$, then $f(3) + f(2) = \cdots$

(a) 125

(b) 25

(c) 150

(d) 100

(9) If \triangle ABC, in which m (\angle A) = 60°, m (\angle C) = 40°, c = 8.4 cm., then a \simeq

(a) 5.3

(b) 11.3

(c) 22.6

(d) 12

(10) If \triangle ABC, which $b = 2 r \times \cdots$

(a) sin A

(b) sin B

(c) cos B

(d) sin C

(11) In \triangle ABC, if $2 \sin A = 3 \sin B = 4 \sin C$, then $a : b : c = \cdots$

(a) 2:3:4

(b) 4:3:2

(c) 3:2:4

(d) 6:4:3

(12) In \triangle ABC which a = b = 5 cm., c = 6 cm., then $\cos A = \cdots$

(a) 0.4

(b) 0.6

(c) 0.8

(d) 0.2

(13) $\log x - \log 3 = \log 9$, then $x = \cdots$

(a) 3

(b) 4

(c) 9

(d) 27

(14) $\log 2 = X$, $\log 3 = y$, then $\log 6 = \dots$

(a) X + y

(b) χ y

(c) X - y

(d) $\log x + \log y$

(15) $\lim_{x \to \infty} (5 x^{70} + 8 x^{30} + 4) = \cdots$

(a) 0

(c) 19

(d) ∞

(16) $\log_3 2 = A \cdot \log_5 3 = B \cdot \text{then } A \times B = \dots$

(a) $\log_5 2$

(b) $\log_2 5$ (c) $\log_3 10$

(d) log 5

(17) $\lim_{x \longrightarrow 4} \left(2x + \sqrt{x} \right)$

(a) 4

(b) 6

(c) 8

(d) 10

$$(18)\sqrt{5} \times \sqrt{2} = \sqrt[6]{x}$$
, then $x = \dots$

(a) 500

(b) 108

(c)72

(d) 1000

(19) If
$$\frac{3^{x} + 2^{x} + 1}{5^{x} + 10^{x} + 15^{x}} = \frac{1}{25}$$
, then $x = \dots$

(a) 1

(b) 2

(c) - 1

(d) - 2

(20) Measure of greatest angle in the triangle whose sides length 7 cm. , 5 cm.

, 3 cm. =

(a) 120°

(b) 150°

(c) 60°

(d) 30°

(21) Length of diameter in the circumcircle of the triangle ABC which a = 8 sin A is unit.

(a) 4

(b) 5

(c) 8

(d) 8 sin A

(22) The solution set of the equation : $4^{x} + 2^{x+1} = 8$ in \mathbb{R} is

(a) $\{1\}$

(b) $\{-1\}$

(c) $\{-2\}$

(d) Ø

(23) $\lim_{x \to 5} (3 + x) = \dots$

(a) 8

(b) 2

(c) 4

(d) - 2

(24) $\lim_{x \to 2} \frac{x^2 - 4}{x - 2} = \dots$

(a) 2

(b) - 4

(c) 4

(d) undefined.

(25) $\lim_{x \to 3} \frac{\sqrt{x+1}-2}{x-3} = \dots$

(a) $\frac{1}{4}$

(b) $\frac{1}{8}$

(c) 12

(d) 3

(26) $\lim_{x \to \infty} \frac{9 x + 5}{3 x - 7} = \dots$

(a) 3

(b) 6

(c) 9

(d) ∞

(27) $\lim_{x \to 1} \frac{x^3 - x^2}{x^3 + 1} = \dots$

(a) 0

(b) - 1

(c) not exist.

(d) $\frac{1}{3}$

Second Essay questions

Answer the following questions :

Represent the function f graphically, then show the range and determine its type whether it is even, odd or otherwise where: $f(X) = (X - 2)^3 + 1$

2 Find:
$$\lim_{x \to 3} \frac{x^5 - 243}{x - 3}$$

Assiut Governorate



Kousya Directorate

First

Multiple choice questions

Choose the correct answer from the given ones:

(1) The domain of the function $f: f(X)$:	$= \frac{2 x + 1}{x - 2} $ is
--	-------------------------------

(a) IR

(b) $\mathbb{R} - \left\{ -\frac{1}{2} \right\}$ (c) $\mathbb{R} - \left\{ -\frac{1}{2}, 2 \right\}$ (d) $\mathbb{R} - \left\{ 2 \right\}$

(2) If $\chi^{\frac{3}{2}} = 64$, then $\chi = \dots$

(a) 512

(d) 2

(3) The type of the function $f: f(X) = \frac{\sin X}{x}$ is

(a) even.

(b) odd.

(c) neither even nor odd.

(d) linear.

(4) $\lim_{x \to 0} (2x^2 + 3) = \cdots$

(a) 3

(b) 2

(c) 7

(d) 5

(5) The range of the function f: f(X) = |X| is

(a) [0,∞[

(b) $]0, \infty[$ (c) $]-\infty, 0]$ (d) $]-\infty, 0[$

(6) $\lim_{x \to 0} \frac{x^2 - x}{x} = \dots$

(a) zero

(b) 1

(c) - 1

(d) does not exit.

(7) The vertex point of the cuve of the function f: f(x) = |x+3| - 2 is

(a)(3,2)

(b) (-3, -2) (c) (-3, 2) (d) (3, -2)

(8) The side length of an equilateral triangle is 9 cm. , then the area of its circumcircle equals cm.2

(a) 9 TT

(b) 27π

(c) 81 π

(d) 72π

(9) The point of symmetry of the curve of the function $f: f(x) = \frac{1}{x-3} + 4$ is

(a) (3, -4)

(b) (-3, -4) (c) (3, 4)

(d) (-3,4)

(10) $\lim_{x \to 2} \frac{x^3 - 8}{x - 2} = \dots$

(a) 1

(b) 12

(c) 0

(d) 3

(11) The solution set of the inequality $|2 \times +3| \le 1$ is

(a) Ø

(b) [-2,-1] (c)]-2,-1[

(b) ab cos C


(c) cos C

(a) cos A


(d) 2 ab cos C

(26) In the opposite figure:

The graph of function f



- (a) 2
- (b) 3
- (c) 1
- (d) not exist.



- (27) By solving the triangle ABC in which a = 5 cm. , b = 7 cm. , $m (\angle C) = 65^{\circ}$, then $c \simeq \cdots cm$.
 - (a) 4.4

- (b) 2.1
- (c) 6.7
- (d) 8.2

Second Essay questions

Answer the following questions:

- Draw the graph of the function $f: f(X) = \begin{cases} X^2, & X < 0 \\ X, & X \ge 0 \end{cases}$ and deduce from the graph its range and its type of being odd, even of otherwise.
- Find the S.S. of the equation: $\log_{\chi} 81 = 4$

Qena Governorate



Mathematics Inspection

First Multiple choice questions

Choose the correct answer from the given ones:

- (1) Solution set of inequality: $|X+2| \le 3$ in \mathbb{R} is
 - (a) R

- (b) Ø
- (c) [-5,1] (d)]-5,1[
- (2) In \triangle ABC if a = 4 cm., $m(\angle A) = 30^{\circ}$, then length radius of circumcircle \triangle ABC = cm.
 - (a) 4

- (b) 2
- (c) 12
- (d) 24

(3) Domain of function $f: f(x) = \sqrt[3]{x-2}$ is			
(a) $[-2,\infty[$	(b) $[-2,2]$	(c) $]-\infty$, 2]	(d) ℝ
(4) $\lim_{x \to \infty} \left(\frac{x}{2 - x^2} \right) = \cdots$	50,000a		
(a) 0	(b) $\frac{1}{2}$	(c) 2	(d) - 1
(5) If $\log 2 = X$, $\log 3 = y$, then $\log 6 = \cdots$		
(a) X y	(b) $X \div y$	(c) $X + y$	(d) X^y
(6) $\lim_{x \to -3} \left(\frac{x^2 + 4x + 3}{x^2 - 9} \right) =$	anatomie		
(a) 2	(b) 0	(c) $\frac{-1}{3}$	(d) $\frac{1}{3}$
(7) In \triangle ABC if $a = 5$ cm., then $c = \cdots \cdots cm$.	$m (\angle B) = 120^{\circ} \text{ and}$	d surface area of Δ Al	$BC = 10\sqrt{3} \text{ cm}$
(a) 5	(b) 8	(c) 7.2	(d) 10
(8) If $5^{x} = 2$, then $5^{x+2} = 4$	940000 000 0000		
(a) 5	(b) 2	(c) 25	(d) 50
(9) In \triangle XYZ if $X = 5$ cm.	y = 7 cm, $z = 8$	3 cm., then $m (\angle Y)$	=
(a) 30°	(b) 60°	(c) 45°	(d) 120°
(10) Solution set of equation :	$\chi^{\frac{7}{2}} = 128$ in \mathbb{R} is	ararrer.	
(a) {4}	(b) {7}	(c) {2}	$(d) \{\pm 4\}$
(11) $\lim_{X \to 0} \left(\frac{X^2 + 2X}{X} \right) = \cdots$			
(a) - 2	(b) 2	(c) 0	(d) 1
(12) Point vertex curve of fun			
(a) $(-2, 1)$	(b) (2, 1)	(c) $(-2, -1)$	(d) $(2,-1)$
$(13) \log_2 6 \times \log_6 2 = \cdots$	no.		
(a) 0	(b) 2	(c) 1	(d) 3
(14) In \triangle ABC if : 3 sin A = 6			
(a) 3:4:6		(c) 3:6:4	
(15) The curve of function $f: f(x) = \frac{1}{x+1}$ symmetry around point			
(a) $(-1,0)$	(b) (0 , 1)		(d) $(1,0)$
(16) $\lim_{x \to 1} \frac{x^4 - 1}{x^2 - 1} = \dots$			
(a) 6	(b) 3	(c) 2	(d) 1

(17) If $\log x \in]0$, 1[, then $x \in ...$

(a)
$$]1,2[$$

(b)
$$]0,1[$$
 (c) $]1,10[$ (d) $]1,\infty[$

(18) $\lim_{x \to 1} (10) = \cdots$

$$(c) - 10$$

(19) In \triangle DEF if , m (\angle E) = 35° , m (\angle F) = 40° , EF = 12 cm. , then ED \simeq to nearest centimeter.

$$(d)\sqrt{3}$$

(20) In \triangle ABC if AB = 3 cm., BC = 5 cm., m (\angle B) = 120°, then AC = cm.

(b)7

(21) Which of following functions represent an even function

(a)
$$f(X) = X^4 + 1$$

(b)
$$f(X) = X^3 + 1$$

(b)
$$f(X) = X^3 + 1$$
 (c) $f(X) = X \cos X$ (d) $f(X) = X^2 \sin X$

(22) $\lim_{n \to \infty} \frac{5 x^{-3} + x^{-1} + 5}{2 x^{-3} + 2 x^{-1} + 7} = \dots$

(a)
$$\frac{5}{7}$$

(b)
$$\frac{7}{5}$$

(c)
$$\frac{1}{2}$$
 (d) $\frac{5}{2}$

(23) Function f where $f(X) = a^X$ is increase on its domain when

(a)
$$a = 1$$

(b)
$$a > 1$$

(c)
$$a = -1$$

(c)
$$a = -1$$
 (d) $0 < a < 1$

(24) $\lim_{x \to 2} \frac{2x^2 - x - k}{x^2 - x - 2} = \frac{7}{3}$, then $k = \dots$

$$(c) - 6$$

(d) 6

(25) $\lim_{x \to 0} \frac{(x+2)^5 - 32}{x} = \cdots$

(b) 16

(c) 32

(d) 80

(26) If $\log_4 (X + 1) = 1$, then $X = \dots$

(27) If $f(X) = 3^X$, then solution set of equation: f(X+1) - f(X-1) = 24 is

(a) $\{2\}$

(b) $\{3\}$

(c) $\{8\}$ (d) $\{0\}$

Essay questions Second

Answer the following questions:


Represent graphically function $f: f(x) = (x-1)^3 + 2$ from drawing determine the range of the function and discuss its monotonicity.

Find value of: $\lim_{x \to 0} \frac{\sqrt{1+x}-1}{x}$



ကြောင်္ကျာပိုက်မျှာတွင်ပြည်တွင်ပြည်လျှင်





~ 8°

Everage

اوتحانات رقور (2)







FIRST

Examinations of some governorate's schools

Cairo Governorate



Mathematics Department Futures Language Schools

First

Multiple choice questions

Choose the correct answer from the given ones:

(1) If f is an odd function, a \subseteq the domain of f, then $f(a) + f(-a) = \cdots$
--

(a) 2 (a)

- (b) 2(-a)
- (c) zero
- (d) (a)

(2) If
$$f(x) = \sqrt{x+4}$$
, $g(x) = \sqrt{6-x}$, then $(f+g)(5) = \cdots$

- (a) undefined
- (b) zero
- (d) 4

(3) The domain of the function
$$f: f(x) = \begin{cases} x+3 & x>3 \\ 6 & x<3 \end{cases}$$
 is

(a) $\{3\}$

- (b) $\mathbb{R} \{3\}$ (c) $[3, \infty[$
- (d) R

(4) If y = f(X) is a real function, then its image by translation 3 units right is $g(X) = \dots$

- (a) f(X-3)
- (b) f(x+3)
- (c) f(x) + 3
- (d) f(x) 3

(5) The solution set of the equation: $\log_5 y = 2$ in \mathbb{R} is

(a) $\{25\}$

- (b) $\{25,625\}$ (c) $\{\frac{1}{25},625\}$ (d) $\{125,625\}$

(6) If $\log_2 x = 3$, then $\log_x 2 = \cdots$

(a) 2

- (b) $\frac{1}{3}$
- (c) 8

(7) If $5^{X+1} = 7^{X+1}$, then $3^{X+1} = \cdots$

(a) zero

- (b) 3
- (c) 2
- (d) 1

(8) Solution set of the equation : $\log_{\mathcal{X}}(X+6) = 2$ in \mathbb{R} is

- (a) $\{3, -2\}$
- (b) $\{3\}$
- (c) $\{3,1\}$ (d) $\{6,1\}$

(9) The domain of the function: $f(x) = \frac{1}{x^2 - 4}$ is

- (a) $\{2, -2\}$

- (b) [2,-2] (c) $\mathbb{R} [2,-2]$ (d) $\mathbb{R} \{2,-2\}$

(10) The solution set of the inequality: |x-3| < 6 is

- (a)]-1,7[
- (b) $\mathbb{R} [-3, 9]$ (c) $\mathbb{R} [-1, 7]$ (d)]-3, 9[

(11) If $2^{X+1} = 8$, then $X = \dots$

- (b) 2
- (c) 3

(a) a = 1

- (b) a > 1
- (c) 0 < a < 1 (d) a = -1

(13) If $3^a = 4$, then $9^a = \cdots$	MARIO AND S		
(a) 7	(b) 12	(c) 16	(d) 25
(14) The equation of axis of s	ymmetry of the cur	ve of the function .	f: f(X) = X + 3 - 2
is			
(a) $X = -3$		(c) $y = -3$	(d) $y = -2$
(15) $\lim_{x \to 0} \frac{\sqrt{x+1}-1}{x} = \cdots$			
2	(b) 2	(c) zero	(d) 1
(16) $\lim_{x \to \infty} x^{-5} = \cdots$			
(a) ∞	(b) - 5	(c) 5	(d) zero
(17) $\lim_{x \to 1} \frac{x^2 - x}{x^3 - 1} = \dots$			
(a) zero	(b) $-\frac{1}{3}$	(c) $\frac{1}{3}$	(d) does not exist
(18) $\lim_{X \to \infty} \frac{X^{-3} + 3X^{-2} + 1}{X^{-2} + X^{-1} + 3}$	=		
(a) 2	(b) 1	(c) 3	(d) $\frac{1}{3}$
(19) $\lim_{x \to 3} \frac{(x-6)^2 - 9}{x^2 - 9} = \cdots$			
(a) - 1	(b) 3	(c) 1	(d) 2
(20) $\lim_{x \to 1} \frac{x^2 + 5x - 6}{x^2 - 1} = \cdots$			
(a) 1	(b) 5	(c) 6	(d) 3.5
(21) In \triangle ABC, $b^2 + c^2 - a^2$	$^2 = 2 \text{ bc} \times \cdots$		
(a) cos A		(c) cos C	(d) cos B
(22) In triangle ABC If: $\frac{a}{\sin A}$	$\frac{1}{4} = 6 \text{ cm.}$, then th	e circumference	of the circumcircle of
(a) 12π	(b) 6 π	(c) 5 π	(d) 9 π
(23) In \triangle ABC, $c = 7$ cm.,	$m (\angle A) = 70^{\circ}$	$m (\angle B) = 40,$	then $b \approx \cdots cm$.
(a) 3.7	(b) 4.8		
(24) If ABC is a triangle in w then the measure of the			c = 8 cm.
(a) 60°	(b) 30°	(c) 90°	(d) 120°
(25) The diameter length of t is $4\sqrt{3}$ cm. equals		n an equilateral tri	angle whose side length
(a) 8	(b) $4\sqrt{3}$	(c) 4	(d) $2\sqrt{3}$

- (26) In \triangle ABC, if $2 \sin A = 3 \sin B = 4 \sin C$, then a: b: c =
 - (a) 6:4:3
- (b) 4:3:2 (c) 3:4:6 (d) 2:3:4
- (27) In \triangle ABC, if m (\angle B) = 60°, m (\angle C) = 30°, c = 4 cm., then b = cm.
 - (a) 4

- (28) ABCD is a parallelogram in which: AB = 9 cm. BC = 13 cm. AC = 20 cm. then the length of BD equals cm.
 - (a) 10

- (b) 5
- (c) 18.5
- (d) 20

Essay questions Second

Answer the following questions:

- 1 Find: $\lim_{x \to 1} \frac{x^3 2x + 1}{x^2 + x 2}$
- Write the steps to find: $\lim_{x \to 3} \frac{\sqrt{x+1}-2}{x-3}$
- If $f(x) = x^2 1$ graph the function showing its domain, range, and monotony
- If $f(x) = 7^x$, then find the value of x satisfying $f(2x-1) + f(2x+1) = \frac{50}{40}$

Cairo Governorate



Multiple choice questions First

Choose the correct answer from the given ones:

- - (a) 2

- (2) The point of symmetry of the function $f: f(x) = \frac{2x-1}{x}$ is
 - (a)(2,0)

- (b) (0, 2)
- (c) (-2,0) (d) (2,-1)
- (3) The solution set of $|2 \times -3| \le 3$ in \mathbb{R} is
 - (a) [-3,3]
- (b)]-3,3[(c) [0,6] (d) [0,3]

- (4) $\lim_{h \to 0} \frac{(x+h)^7 x^7}{h} = \dots$

- (b) $6x^7$

- (5) In \triangle ABC, if $b^2 = c^2 + a^2 ac$, then m (\angle B) =
 - (a) 30°


- (b) 45°
- (c) 60°
- (d) 120°

- (6) $5^{X-3} = 4^{3-X}$, then $X = \dots$
 - (a) 3

- (b) 4
- (d)0

- (7) The range of the function $f: f(X) = \begin{cases} 1, & X \le 0 \\ 0, & X > 0 \end{cases}$
 - (a) R

- (b) $\mathbb{R} \{0, 1\}$ (c) $\{0, 1\}$
- (d) $\mathbb{R} [0, 1]$
- (8) The opposite figure represents $f: f(X) = \cdots$
 - $(a)\frac{1}{2}$
 - $(b)-\frac{1}{\gamma}$
 - $(c)\frac{1}{|x|}$
 - $(d)\frac{1}{x} + 5$



- (9) In \triangle XYZ, X = 5 cm., y = 7 cm., $m (\angle Z) = 65^{\circ}$, then $z \simeq \cdots cm$.
 - (a) 5.7

- (b) 6.7
- (c) 7.5
- (d) 44

- (10) If $\log_{x} 2 = 3$, then $\log_{2} X = \cdots$
 - (a) X

- (b) 1
- (c) $\frac{1}{3}$
- (d)3

- (11) The solution set of |X| + 3 = 1 in \mathbb{R} is
 - (a) $\{2\}$

- (b) $\{-2\}$ (c) $\{-2, 2\}$

- (12) $\lim_{y \to 2} \frac{y^5 32}{y^2 4} = \dots$

- (b) 40
- (c) 60
- (d) 80
- (13) If the curve of the function $f: f(X) = \log_4 (1 aX)$ passes through the point $(\frac{1}{8}, -\frac{1}{2})$, then a =
 - (a) 1

- (b) 2
- (c) 3
- (14) The domain of the function $f: f(x) = \sqrt{9-x}$ is
 - (a) [9,∞[

- (b) $]-\infty, 9]$ (c) $[-9, \infty[$
- (15) In $\triangle XYZ$, $\sin X = 2 \sin Z$ and YZ = 6 cm., then $XY = \cdots$

- (d) 12

- (a) 5 (16) $\lim_{x \to \infty} \frac{6x^2 3x + 6}{1 + 4x + 2x^2} = \dots$

- (b) 3
- (c) 4
- (17) If f is an odd function and a \subseteq its domain, then $f(a) + f(-a) = \cdots$
 - (a) 0

- (b) 2
- (c) 2 f (a)
- (d) f(a)

- (18) $f(X) = 2^{1-X}$, then $f(-1) = \cdots$
 - (a) 0

- (c) 2
- (d) 4

- (19) $\lim_{x \to 3} \frac{\sqrt{x+1}-2}{x-3} = \dots$

- (d) does not exist.

f(x)

- (20) $f(X) = 3^{X+1}$, then $f(X+1) \times f(-X) = \cdots$

- (b) 3
- (c) 9
- (d) 27
- (21) The side length of an equilateral triangle is 9 cm., then the area of its circumcircle equals cm².
 - (a) 9 T

- (b) 27 TT
- (c) 81 T
- (d) 72π
- (22) The opposite figure represents the curve of the function f, then $\lim_{x \to 0} f(x) = \cdots$
 - (a) 1
 - (b) 4
 - (c) 0
 - (d) does not exist.
- (23) If $\lim_{x \to 2} \frac{a x}{3} = 6$, then $a = \dots$

- (b) 4
- (c) 6
- (d) 9

- (24) In \triangle ABC, if b = c, then $\cos C = \dots$

- (b) $\frac{a}{b}$

- (25) If $\log_2 x = 5$, then the exponential form of it is
 - (a) $\chi^2 = 5$
- (b) $2^5 = X$ (c) $5^2 = X$
- (d) $x^5 = 2$
- (26) In \triangle ABC, if m (\angle A) = 110°, m (\angle B) = 34°, c = 19 cm. , then b to nearest cm. = cm.
 - (a) 14

- (c) 19.8
- (d) 30.4

- (27) $\lim_{x \to \infty} \frac{6}{3 x^2} + \frac{8 x}{2 + x} = \cdots$

- (b) 4
- (d) 8
- (28) If $\log 3 = x$ and $\log 4 = y$, then $\log 12 = \dots$
 - (a) X y
- (b) X + y
- (c) X y

Second Essay questions

Answer the following questions:

Find: $\lim_{x \to 2} \frac{x^2 - 5x + 6}{x^2 - 4}$

- Graph the function $f: f(X) = \begin{cases} |X|, & X \le 0 \\ X^3, & X > 0 \end{cases}$, then
 - (1) deduce its range.

(2) discuss the monotony.

- Find: $\lim_{x \to 1} \frac{(x+1)^5 32}{x-1}$
- Find the solution set of: $3^{x+2} 3^{x+1} = 18$ in \mathbb{R}

Cairo Governorate



Ain Shams Educational directorate

First Multiple choice questions

Choose the correct answer from the given ones:

- (1) The type of the function $f: f(x) = \frac{\sin x}{x}$ is
 - (a) even.

(b) odd.

(c) neither even nor odd.

- (d) linear.
- (2) The domain of the function $f: f(X) = \frac{2X+1}{X-2}$ is
 - (a) IR

- (b) $\mathbb{R} \left\{ -\frac{1}{2} \right\}$ (c) $\mathbb{R} \left\{ -\frac{1}{2}, 2 \right\}$ (d) $\mathbb{R} \left\{ 2 \right\}$

- (3) 1 + log 2 =
 - (a) log 5

- (b) log 2
- (c) log 20
- $(d) \log 5$

- (4) In \triangle XYZ, $y^2 + z^2 x^2 = 2$ y z ×
 - (a) cos X
- (b) sin Z
- (c) cos Z
- (d) sin X

- (5) $\lim_{x \to \infty} \frac{3x}{4x+5} = \cdots$
 - (a) ∞

- (b) $\frac{3}{4}$
- (c) $\frac{1}{5}$
- (d) zero
- (6) The set of the real roots of the equation $(x-2)^4 = 16$ equals
 - (a) $\{0\}$

- (b) $\{4\}$
- (c) {8}
- $(d) \{0,4\}$
- (7) The range of the function f: f(x) = |x-2| is
 - (a) $-\infty$, 2

- (b) $\begin{bmatrix} -2, \infty \end{bmatrix}$ (c) $\begin{bmatrix} 0, \infty \end{bmatrix}$ (d) $\begin{bmatrix} 2, \infty \end{bmatrix}$
- (8) If $\log x \log 2 = \log 4$, then $x = \dots$
 - (a) 4

- (b) 6
- (c) 8
- (d) 16

- (9) $\lim_{x \to 2} \frac{x^5 32}{x^3 2^3} = \dots$

- (b) $\frac{5}{2}$
- (c) zero
- (d) $6\frac{2}{3}$

(10) In \triangle ABC, $\frac{a}{\sin A} = 6$,	then the length of the	diameter of its circu	mcircle
is length unit	s		
(a) 6	(b) 12	(c) 3	(d) 9
(11) If $\sqrt[3]{x^2} = 9$, then $x \in$			
(a) {27}	(b) $\{27, -27\}$	(c) $\{1\}$	(d) Ø
(12) If $\log_3 x = 2$, then x:	=		
(a) 3	(b) 5	(c) 8	(d) 9
(13) $\lim_{x \to -1} 3 x^2 = \dots$			
(a) 2	(b) 3	(c) 4	(d) 5
(14) In \triangle ABC, $\cos (A + B)$			
(a) $\frac{a^2 + b^2 - c^2}{2 ab}$	(b) $\frac{a^2 + c^2 - b^2}{2 ab}$	(c) $\frac{b^2 + c^2 - a^2}{2 bc}$	(d) $\frac{c^2 - a^2 - b^2}{2 ab}$
(15) If $2^{X+1} = 8$, then $X =$			
(a) 1	(b) 2	(c) 3	(d) 4
(16) The solution set of the	inequality: $ 2 X + 3 $	≤ 1 in ℝ is	
(a) IR		(c) $[-2, -1]$	(d) Ø
(17) The vertex of the curve	of the function f whe	ere $f(x) = (1 + x)^2$	– 3 is
	(b) $(1, -3)$		
(18) $\lim_{x \to 4} \frac{(x-3)^2 - 1}{x-4} = \cdots$			
(a) zero	(b) 2	(c) 3	(d) 4
(19) $\lim_{x \to \infty} \frac{2x}{\sqrt{9x^2 + 1}} = \cdots$			
(a) $\frac{2}{9}$	(b) zero	(c) $\frac{2}{3}$	(d) ∞
(20) If the perimeter of trian	gle ABC equals 15 cn	$m. , m (\angle A) = 82^{\circ}$	$m (\angle B) = 47^{\circ}$
, then the length of AB	≃ cm.		
(a) 6	(b) 7	(c) 5	(d) 8
(21) In \triangle ABC, m (\angle A) =	45°, the length of the	radius of its circumo	circle = 6 cm.
, then $a = \cdots \cdots cm$			
(a) 13	(b) $6\sqrt{2}$	(c) 12	$(d)\sqrt{2}$
(22) If $y = f(X)$ is a real furupwards is $g(X) = \cdots$		by translation 3 unit	s vertically
(a) $f(X-3)$	(b) $f(X + 3)$	(c) $f(x) + 3$	(d) $f(x) - 3$

(23)
$$\lim_{X \to -2} \frac{3 x^2 - 12}{X + 2} = \dots$$

(b) - 3

(c) 12

(d) - 12

(24) DEF is a triangle in which m (\angle D) = 80° and m (\angle E) = 60°, if f = 12 cm. , then $d = \cdots cm$.

(a) $\frac{12 \sin 80^{\circ}}{\sin 40^{\circ}}$

(b) $\frac{12 \sin 80^{\circ}}{\sin 60^{\circ}}$ (c) $\frac{12 \sin 40^{\circ}}{\sin 80^{\circ}}$ (d) $\frac{12 \cos 80^{\circ}}{\cos 40^{\circ}}$

(a) ∠ A

(b) ∠ B

(c) \(C

(d) right.

(26) In \triangle XYZ, x = 5 cm., y = 3 cm., $m (\angle Z) = 120^{\circ}$, then $z = \cdots$ cm.

(a) 7

(b) 6 (c) $3\sqrt{3}$

(d) 4

(27) The solution set of the equation : |X-2|=3 is

(a) $\{2,3\}$

(b) $\{-1,5\}$ (c) [-1,5] (d) $\{5,-5\}$

(28) If $3^{x} = 5$, then $x = \cdots$

(a) 3

(b) $\log_3 5$ (c) $\log_5 3$ (d) $\frac{5}{3}$

Second Essay questions

Answer the following questions:

- Find the solution set in $\mathbb{R}: |X-3| = |X+1|$
- If $f(x) = 5^x$, find the value of: $\frac{f(x+4) f(x+3)}{f(x+5) f(x+4)}$
- 3 Find: $\lim_{x \to \infty} \left(\frac{x}{2x+1} + \frac{3x^2}{(x-2)^2} \right)$
- 4 Find: $\lim_{x \to -1} \frac{x+1}{\sqrt{x+5}-2}$

Giza Governorate



Awseem Educational Directorate

Multiple choice questions First

Choose the correct answer from the given ones:

(1) The type of the functions $f: f(X) = X^2$ where $f: \mathbb{Z}^+ \longrightarrow \mathbb{Z}$ is

(a) even.

(b) odd.

(c) neither even nor odd.

(d) constant.

(2) If $f: f(x) = 2$, then	nen the range of the fu	nction f is	***
(a) R	(b) IR +	(c) {2}	$(d)\mathbb{R}-\{2\}$
(3) The range of the fu	unction $f: f(X) = \frac{1}{X} + 1$	is	
(a) R	(b) ℝ − {2}		(d) $\mathbb{R} - \{1\}$
(4) The S.S. of the equ	nation: $ X - 2 + 1 = 0$	is	
(a) R	(b) ∅	$(c){3}$	(d) $\{-1\}$
(5) The axis of symme	etry of the function $f: f$	$(X) = 2 - (X - 1)^2$ i	s $x = \cdots$
(a) 1	(b) - 1	(c) 2	(d) 3
(6) If $5^{X+2} = 125$, th	en $X = \cdots$		
(a) 2	(b) 1	(c) 3	(d) 4
(7) If $9 \times 3^{2-x} = 81^-$	1 , then $X = \cdots$		
(a) 6	(b) 7	(c)8	(d) 9
(a) 6 (8) $((2)^7 \div (2)^5)^{\frac{1}{2}} = \cdots$			
(a) 2	(b) - 2	$(c)\frac{1}{2}$	$(d)\frac{-1}{2}$
(9) The two curves of at $x = \cdots$	the two functions $f: f$ (.	$(x) = 2^{x}, g : g(x)$	= 3 ^x will intersect
(a) 2	(b) 3	(c) zero	(d) 5
(10) If $\log_{x} (5 x) = 2$,	then $x \in \{\dots \}$		
(a) $\{0,5\}$	(b) {5}	(c) {0}	(d) $\{2\}$
$(11)\log_8\log_2\log_3(X$	-4) = $\frac{1}{3}$, then $x = \cdots$		
(a) 8	(b) 48	(c) 90	(d) 85
$(12) \log 125 - \log 6 + \log 6$	og 48 = ·····		
(a) 3	(b) 6	(c) 7	(d) 8
(13) If $\log_2 X + \log_4 X$	$= 3$, then $X = \cdots$		
(a) 2	(b) 3	(c) 4	(d) 5
(14) If $2^{x} = 7$, then x	~ ··········		
(a) 2.25	(b) 2.81	(c) 2.85	(d) 3
(15) $\lim_{x \to 3} \frac{x^3 - 27}{x^2 - 2x - 3}$	-=		
(a) 6.5	(b) 6.75	(c) 7	(d) 7.5
(a) 6.5 (16) $\lim_{x \to 2} (10) = \dots$	eatra -		
(a) 2	(b) 5	(c) 10	(d) 8

(17)
$$\lim_{x \to 2} \frac{x^4 - k^4}{x - k} = 32$$
, then $k = \dots$
(a) zero (b) 1
(18) $\lim_{x \to \infty} (3x^{-5} + 4x^{-2} + 5) = \dots$
(a) 5 (b) ∞

(19) $\lim_{X \to \infty} X^{-4} = \dots$

(a) zero

(c) 2

(c) 12

(d) ∞

(d) zero

(d)3

(20) In \triangle XYZ, the expression $\frac{\chi^2 + y^2 - z^2}{2 \chi y} = \cdots$

(a) cos X

(c) cos Z

(d) sin Y

(21) $\lim_{h \to 0} \frac{(2+3h)^5-32}{2h} = \cdots$

(a) 32

(c) 80

(d) 120

(22) In triangle ABC if $a^2 = b^2 + c^2 + bc$, then $m (\angle A) = \cdots$

(a) 120

(b) 60

(c) 45

(d) 30

(23) In triangle ABC if m (\angle C) = 30°, AB = 14 cm., then the circumference of the circle = cm.

(a) 28π

(b) 30 π

(c) 28

(d) 335 T

(24) The radius length of the circumcircle of triangle ABC in which m (\angle A) = 30° , a = 10 cm. is

(25) If r is the radius of the circumcircle of triangle XYZ, then $\frac{2 \text{ y}}{\sin x} = \frac{(d) 40}{\sin x}$

(26) ABC is a triangle in which $\cos B = \frac{c}{2a}$, then the triangle will be

(a) scalene.

(b) right-angled. (c) isosceles. (d) equilateral.

(27) In triangle XYZ: $y^2 + z^2 - x^2 = 2$ y z

(a) cos X

(b) sin Z

(c) cos Z

(28) In triangle XYZ if: $2 \sin X = 3 \sin Y = 4 \sin Z$, then $X: y: z = \cdots$

(a) 2:3:4

(b) 6:4:3 (c) 3:4:6 (d) 4:3:2

Essay questions Second

Answer the following questions:

11 Draw the graph of the function $f: f(x) = x^3 + 1$ and deduce from the graph its range and its monotony.

- Find in \mathbb{R} the S.S. of the equation: $5^{x+1} + 5^{x-1} = 26$
- 3 Find : (1) $\lim_{x \to 1} \frac{x^3 2x + 1}{x^2}$
- (2) $\lim_{x \to 0} \frac{(x+1)^{11}-1}{x}$
- Find the value of : (1) $\lim_{x \to 3} \frac{x^2 8x + 15}{x 3}$ (2) $\lim_{x \to \infty} (x^5 + x^2 1)$
 - Giza Governorate



Mathematics inspection


Multiple choice questions **First**

Choose the correct answer from the given ones:

- (1) The opposite figure represents a function it's range is
 - (a) $\{1\}$

(b) $\{1, -1\}$

(c) $\{-1\}$



- (2) $\lim_{x \to \infty} \frac{2x+3}{5x^2+4} = \dots$
 - (a) 2

- (b) zero
- (c) $\frac{3}{4}$
- (d) $\frac{4}{10}$
- (3) ABC is an equilateral triangle, its side length is $5\sqrt{3}$ cm., then the length of the diameter of circumcircle = cm.
 - (a) 5

- (b) 10
- (c) 15
- (d) 20
- (4) The exponential function $f: f(X) = a^X$ is increasing when
 - (a) a > 0

- (b) a > 1 (c) a = 1
- (d) 1 > a > 0

- (5) $\lim_{x \to 1} \frac{x^7 1}{x 1} = \dots$
 - (a) 35

- (b) 7
- (c) 42
- (6) In \triangle ABC, m (\angle A): m (\angle B): m (\angle C) = 3:5:4, then c^2 : a^2 =
 - (a) $\sqrt{6}:2$

- (b) 2:3
- (c) 4:3
- (7) f is a function, where $f(x) = (x-2)^2$, then the equation of its symmetric axis is $\chi = \cdots$
 - (a) 2

- (b) 2
- (c) 1 (d) 1

(8) $\lim_{x \to 1} \frac{2x+k}{x+1} = 5$, th	en k =		
(a) 2	(b) 5	(c) 8	(d) 10
(9) In \triangle ABC, $\frac{a}{\sin \Delta} = 6$	then the radius length o	of circumcircle = ·····	cm.
(a) 2	(b) 3	(c) 5	(d) 6
(10) $\log_3 5 \times \log_2 3 \times \log_3 5$	g ₅ 16 = ······		
(a) 30		(c) log 10000	(d) log 30 240
(11) The curve of the fund	ection g : g (X) = $X^2 + 4$ is	the same curve of the	ne function
$f: f(X) = X^2$ by tra	nslation of magnitude 4 u	nits in direction of	
A CONTRACTOR OF THE PARTY OF TH		(c) Oy	(d) Oy
(12) $\lim_{x \to 1} \frac{x^2 - x^{-2}}{x - x^{-1}} = $,		
(a) zero	(b) 1	(c) 2	(d) - 2
(13) In \triangle ABC, m (\angle A)	$=30^{\circ}$, $b = 15\sqrt{3}$ cm.	• m (\angle B) = 60°	
, then a =	em.		
(a) 30	(b) 45	(c) 15	(d) 60
(14) S.S. of the equation	$\log_X (X+6) = 2 \text{ in } \mathbb{R} \text{ is } \cdot \cdot$	***********	
(a) $(3, -2)$		(c) $\{3, 1\}$	(d) $\{6,1\}$
(15) $\lim_{x \to 1} (2x-5) = \cdots$	**********		
(a) 2	(b) - 3	(c) 7	(d) zero
(16) In \triangle ABC, $a^2 + b^2$	$-c^2 = \cdots$		
(a) cos A	(b) a b cos C	(c) cos C	(d) 2 a b cos C
(17) The solution set of i	nequality: $ x-2 < 5$ is		
(a) $[-3, 7]$	(b) $]-3,7[$	(c) $\mathbb{R} - [-3, 7]$	(d) $\mathbb{R} -]-3,7[$
(18) $\lim_{x \to 1} \frac{2x-4}{x-2} = \cdots$			
(a) 1	(b) 2	(c) - 2	(d) zero
(19) In \triangle ABC , if sin A	$= 2 \sin C$, BC = 6 cm., t	hen AB =	em.
(a) 2		(c) 4	(d) 6
(20) In \triangle ABC, $\frac{a}{a+b} =$	sin A		
(a) sin B	(b) $\sin A + \sin B$	(c) $\sin A + \sin C$	(d) sin C
(21) In \triangle ABC, $a = 3$ c	m., $b = 5 \text{ cm}$., c	= 7 cm., the measu	ire of the greatest
angle of \triangle ABC is			
(a) 60	(b) 150	(c) 120	(d) 90

(22) If $\chi^{\frac{3}{2}} = 64$, then $\chi = \dots$

(a) 512

- (b) 16
- (c) 4
- (d) 2

(23) $f(x) = \dots$ is an even function.

(a) $\sin x$

- (b) tan 45°
- (c) $X \cos X$ (d) $X^2 + \tan X$

(24) The function $f: f(x) = \frac{5}{x} + 2$, its range is

(a) R

- (b) $\mathbb{R} \{2\}$ (c) $\{2\}$ (d) $\mathbb{R} \{0\}$

(25) The symmetric point of the function f where $f(x) = \frac{2x-1}{x}$ is

(a) (1 , 2)

- (b) (2, 1)
- (c) (-1, 2) (d) (0, 2)

(26) The point of the vertex of the curve of the function $f: f(x) = (x-2)^2 + 3$ is

(a)(2,3)

- (b) (2, -3) (c) (-2, 3) (d) (-2, -3)

(27) If $\log_4 x = 2$, then the equivalent exponential form is

- (a) $\chi^2 = 4$ (b) $\chi^4 = 2$
- (c) X = 8 (d) $X = 4^2$

(28) If $3^{X-2} = 2^{X-2}$, then $X = \dots$

(a) 3

- (b) 2
- (c) 0
- (d)2

Second Essay questions

Answer the following questions:

Without using calculator find the value of:

 $\log_2 \frac{3}{25} + 5 \log_2 5 + \log_2 27 - \log_2 \frac{125}{12} - \log_2 243$

2 Find: (1) $\lim_{x \to 3} \frac{x^3 - 27}{x^2 - 9}$

- (2) $\lim_{x \to \infty} \frac{4x^2 + 1}{x^2 2}$
- 3 Find: (1) $\lim_{x \to 1} \frac{x^2 + 5x 6}{x^2 + 1}$
- (2) $\lim_{x \to 1} \frac{(x+1)^5 32}{x}$

Graph the curve of the function f where f(x) = |x - 3|, deduce the range and monotony of the function and tell whether it is even, odd or otherwise.

Giza Governorate



El-Omrania Educational directorate

Multiple choice questions First

Choose	the	correct	answer	from	the	given	ones	
CHOOSE	uic	COLLECT	allower	TI OTTI	unc	MINCH	OHEG	

(1	The solution of	of the inequality	2x+3	≤ 7 is ······
----	-----------------	-------------------	------	---------------

$$(a)[-5,2]$$

(b)
$$]-5,2[$$

(c)
$$\mathbb{R} - [-5, 2]$$

(b)
$$]-5,2[$$
 (c) $\mathbb{R}-[-5,2]$ (d) $\mathbb{R}-]-5,2[$

(2) In any \triangle ABC: $\frac{\sin(A+B)}{\sin A} = \cdots$

(a)
$$\frac{a}{a+b}$$

$$(b)\frac{a+b}{a}$$

$$(c)\frac{c}{a}$$

$$(d) - \frac{c}{a}$$

(3) Which of the following functions is even function?

(a)
$$y = X \cos X$$

(b)
$$y = x^2 \sin x$$
 (c) $y = x \sin x$

(c)
$$y = X \sin X$$

(d)
$$y = x^3$$

(4) The solution set of the equation $3 \log_5 (x-2) = 6$ is

(b)
$$\{-27\}$$

(c)
$$\{25\}$$

$$(d)\{7\}$$

(5) ABC is equilateral triangle inscribed in a circle of radius length 10 cm.

, then $AB = \cdots cm$.

(d)
$$5\sqrt{3}$$

(6) $\lim_{x \to \infty} \frac{(2x+1)(3-x)}{(x^2+2)} = \frac{\text{(b) } 10\sqrt{3}}{}$

$$(c) - 2$$

(7) The range of the function f: f(X) = |X-2| + 3 is

(b)
$$[3, \infty[$$
 (c) $]3, \infty[$

$$(d)$$
 $]-\infty,2]$

(8) The solution set of the equation : $\sqrt{x^2 - 10 x + 25} = 10$ is

(a)
$$\{-15, 5\}$$

(b)
$$\{-15, -5\}$$
 (c) $\{15, -5\}$ (d) $\{15, 5\}$

(c)
$$\{15, -5\}$$

$$(d)$$
 {15,5}

(9) $\lim_{x \to 5} \frac{x^2 - 25}{x - 5} = \dots$

$$(b) - 5$$

(b)
$$-5$$
 (c) -10

(10) In \triangle ABC, if $a^2 + b^2 - c^2 = ab$, then m (\angle C) =

$$(a)\frac{\pi}{3}$$

$$(b)\frac{\pi}{6}$$

$$(c)\frac{\pi}{4}$$

$$(d)\frac{\pi}{2}$$

(11) $\lim_{x \to 1} \frac{a x + 3}{x^2 + 1} = 5$, then $a = \dots$

$$(c) - 5$$

(12) The solution set of the equation : $5^{X} + 5^{X+1} = 150$ is

(a)
$$\{-2\}$$

(c)
$$\{-3\}$$

$$(d) \{2, 3\}$$

(13) $\lim_{x \to 2} \frac{x^n - 2^5}{x^2 - 2^m} =$	k , then $m + n + k = \cdots$		
(a) 47	(b) 20	(c) 7	(d) 27
(14) If $\log 3 = X$, then	log 90 =		
(a) 9 X	(b) $2 X + 1$	(c) $X + 1$	(d) $2 X + 10$
(15) The domain of the	function $f: f(x) = \sqrt{x}$	-3 is	
(a) IR		(c) [3,∞[(d) $]-\infty, 3[$
(16) In \triangle ABC, $\frac{a}{\sin A}$ triangle ABC =	$+\frac{c}{\sin C} = 20 \text{ cm.}$, then the	e diameter length of t	the circumcircle of
(a) 5	(b) 10	(c) 20	(d) 40
(17) The axis of symme	etry of the function $f: f$		
(a) $X = 3$		(c) $X + 3 = 0$	
(18) If the domain of th	the function $f: f(X) = \frac{1}{X-1}$	$\frac{1}{a} + 3$ is $\mathbb{R} - \{a\}$, th	nen a ² =
(a) 9		(c) – 4	
(19) In Δ LMN , 3 sin I	$L = 4 \sin M = 5 \sin N$, th	en ℓ : m : n =	de:
(a) 4:5:3	(b) 15:12:20	(c) 20:15:12	(d) 3:4:5
(20) If $5^{x} = 7$, then 5^{3}			
(a) 5 ⁷	(b) 7 ⁵	(c) 12	(d) 35
(21) The number of solu	utions of the \triangle ABC in wh	hich m ($\angle A$) = 112°	a = 7 cm.
b = 4 cm. equal			
(a) 0	(b) 1	(c) 2	(d) 3
(22) $\lim_{x \to 0} \frac{\sqrt{x+9}-3}{5x} =$	= ·····		
	(b) 30		(d) $-\frac{1}{30}$
(23) The domain of the	function $f: f(X) = \log X$	² is	
(a) R*	(b) R	(c) IR -	(d) IR +
	atest angle in triangle who		
(a) 135° 23	(b) 44° 25	(c) 120°	(d) 101° 32
(25) $\lim_{X \to 0} \frac{X}{\cos X} = \cdots$	·············		
(a) zero	(b) 1	(c) not exist	(d) ∞
(26) Δ ABC in which m	$(\angle A) = 80^{\circ}$, m $(\angle C) =$	60° , b = 14 cm., the	en a =
(a) 17.8 cm.	(b) 18.9 cm	(c) 15.6 cm	(d) 21.4 cm

- (27) Which of the following is not a function from X to y?
 - (a) $|y| = 2 x^2$

- (b) $y^3 = 2 X$ (c) y = |X + 1| (d) $y = X^2 + 1$
- (28) $\frac{1}{\log_{a} abc} + \frac{1}{\log_{c} abc} + \frac{1}{\log_{b} abc} = \dots$
 - $(a) (abc)^2$

- (b) abc
- (c) 2 abc
- (d) 1

Second Essay questions

Answer the following questions:

- 11 Draw the curve of the function $f: f(X) = (X-2)^3$ from the graph deduce its range, and discuss its monotony.
- 2 If the volume of a sphere gives by the relation $v = \frac{4}{3} \pi r^3$, if the volume equals 345.45 cm³. Find its radius length.
- 3 Find with steps: $\lim_{x \to 4} \frac{x^3 3x^2 4x}{x \cdot 4}$
- Find with steps: $\lim_{x \to 0} \frac{x^2 + x}{\sqrt{2x + 9} 3}$

Alexandria Governorate



West Education Zone Nabaa Elfekr secondary school


Multiple choice questions **First**

Choose the correct answer from the given ones:

- (1) The domain of the function $f: f(x) = \sqrt[3]{x-5}$ is
 - (a) [5,∞[

- (b) $|5, \infty[$ (c) $|-\infty, 5[$
- (d) R

- (2) The opposite figure represents function.
 - (a) even
 - (b) neither even nor odd
 - (c) odd
 - (d) symmetric about y-axis




- (3) The range of $f: f(X) = \frac{1}{X-2} 1$ is
 - (a) $\mathbb{R} \{2\}$
- (b) $\mathbb{R} \{1\}$ (c) $\mathbb{R} \{-1\}$
- (4) The curve of the function $g: g(x) = (x-2)^2$ is the same of the curve of the function $f: f(X) = X^2$ by translate 2 units in direction of
 - (a) ox

- (b) ox
- (c) oy
- (d) ov

(5) In the opposite figure:

 $\lim_{x \to 2} f(x) = \cdots$

- (a) 1
- (b) zero
- (e) 2
- (d) not exist.



- (6) In \triangle ABC: m (\angle A) = 112°, m (\angle B) = 33°, c = 19 cm., then b \cong cm.
 - (a) 16

- (b) 17
- (c) 18
- (7) The radius of circumcircle of \triangle XYZ when m (\angle X) = 30°, x = 7 cm. equals
 - (a) 10

- (b) 14
- (c) 7
- (d) 21
- (8) The S.S. of the inequality |X| 1 > 0 in \mathbb{R} is
 - (a) $\mathbb{R} [-1, 1]$
- (b)]-1,1[(c) $\mathbb{R}-]-1,1[$ (d) [-1,1]
- (9) The S.S. of the equation: $5^{X+1} = 7^{X+1}$ in \mathbb{R} is
 - (a) $\{1\}$

- (b) $\{-1\}$ (c) $\{zero\}$ (d) $\{5\}$
- (10) The equation of symmetry axis of the function f where $f(x) = (x-2)^2 + 3$ is
 - (a) X = 2

- (b) X = 3

- (11) The S.S. of the equation : $\log_{\mathcal{X}}(X+2) = 2$ is
 - (a) $\{-1\}$
- (b) $\{2\}$
- $(c)\{-1,2\}$

- (12) $\lim_{x \to 0} (2 x^2 + 3) = \cdots$
 - (a) 2

- (b) 3
- (c) 5
- (d) 7
- (13) In \triangle ABC: $\frac{\sin A}{2} = \frac{\sin B}{3} = \frac{\sin C}{5}$, then a: b: c =
 - (a) 4:3:10
- (b) 2:3:5
- (c) 4:6:5
- (d) 4:3:5

- (14) In \triangle ABC: $a^2 + b^2 c^2 = \cdots$
 - (a) cos A

- (b) ab cos C
- (c) cos C
- (d) 2 ab cos C

(15) If f odd function, a	\in domain of f , then f	$(a) + f (-a) = \cdots$	*******
(a) 2 f (a)	(b) 2 f (- a)	(c) zero	(d) f (a)
(16) The point of symme	try of the curve of the fu	unction $f: f(X) = (X)$	$(x-2)^3 + 1$ is
(a) (2, 1)	(b) $(-2,-1)$	(c) $(-2, 1)$	(d) $(2, -1)$
(17) If $2^{X} = 3$, then $X =$			
(a) 2	(b) $\frac{3}{2}$	(c) log ₃ 2	(d) $\log_2 3$
(18) $\lim_{x \to 2} \frac{x^2 - 5x + 4}{x^2 - 1}$	=		
(a) $-\frac{2}{3}$	(b) $\frac{2}{3}$	(c) $\frac{1}{2}$	(d) $-\frac{1}{2}$
(19) $\lim_{x \to \infty} \frac{\sqrt{4x^2 + 1}}{x - 2} =$	************		
(a) 4	(b) 5	$(c) - \frac{5}{2}$	(d) 2
(20) The measure of sma		here $a = 8$ cm., $b = 7$	7 cm. and
its perimeter 21 cm.			
(a) 22° 34	(b) 42° 34		
(21) In \triangle ABC : $\cos A =$	$\frac{2}{5}$, b = 2.5 cm., c = 2	cm. \circ then $a = \cdots$	*****
(a) 2	(b) 2.5	(c) 3	(d) 3.5
(22) The S.S. of the equa	tion $ X-7 = 5$ is		
(a) $\{7, 12\}$	(b) $\{-2,2\}$	(c) $\{7,5\}$	(d) $\{12, 2\}$
(23) The exponential fun	ction which its base (a)	is increasing if	CONTROL I
(a) $a > 0$	(b) $a > 1$	(c) $0 < a < 1$	(d) $a = 1$
(24) If $f(X) = 5^X$, then			
(a) $\{5\}$	(b) $\{2,5\}$	(c) $\{3\}$	(d) $\{2\}$
(25) In \triangle XYZ : X = 5 cr	$m. , Y = 7 cm. , m (\angle Z)$	$=65^{\circ}$, then $Z=\cdots$	
(a) 7.6	(b) 6.7	(c) 7.8	(d) 8.7
(26) In \triangle ABC , if m (\angle	A) = 30° , a = 6 cm., the	$hen \frac{b}{\sin B} = \dots$	
(a) 3	(b) 6	(c) $\frac{1}{2}$	(d) 12
(27) $\lim_{X \to 0} \frac{5 X - 10}{4 X - 8} = \cdots$		-	
(a) $\frac{5}{4}$	(b) zero	(c) 2	(d) $\frac{4}{5}$
(28) If $\lim_{x \to 1} \frac{b}{x+1} = 5$	then b =		
(a) 4	(b) - 1	(c) 1	(d) 10

Second **Essay questions**

Answer the following questions:


- 11 Draw the curve of the function $f: f(x) = (x-2)^2 + 1$, then find its range and monotony and its type.
- Find in \mathbb{R} the solution set of the inequality: $|3 \times -2| \le 7$
- Find: $\lim_{x \to 5} \frac{x-5}{\sqrt{x+4}-3}$
- Find: $\lim_{x \to \infty} \frac{4x^2 + 1}{x^2 2}$



Multiple choice questions First

Choose the correct answer from the given ones:

(1) If $f: \mathbb{R} \longrightarrow \mathbb{R}$ where $f(X) = X^3$, then the figure which represents the function f



- (2) If $5^{X-3} = 4^{3-X}$, then $X = \cdots$

- (c) $\frac{4}{5}$
- (d)0
- (3) The range of the function f where f(x) = |x| is
 - (a) [0,∞[
- (b) $]0, \infty[$ (c) $]-\infty, 0]$ (d) $]-\infty, 0[$


- (4) If $f(x) = 5^x$, then $f(-2) = \cdots$
 - (a) 2

- (b) 5
- (c) $\frac{1}{25}$

- (5) The solution set of the inequality: $|X| 1 > \text{zero in } \mathbb{R}$ is
 - (a) $\mathbb{R} [-1, 1]$
- (b)]-1,1[(c) $\mathbb{R}-]-1,1[$ (d) [-1,1]
- (6) If $4 = \log_2 X$, then the equivalent exponential form is
 - (a) $x^2 = 4$
- (b) $\chi^4 = 2$
- (c) $X = 2^4$ (d) X = 8
- (7) The domain of the function in the figure opposite is



- (b) 0,∞
 - (c)[0,1]
 - (d)]0,2[



- (8) Which of the following functions represents an increasing exponential function on its domain R?
 - (a) $y = 3 (1.05)^{x}$
- (b) $y = 3 \left(\frac{1}{1.05}\right)^{\chi}$ (c) $y = 3 + (0.5)^{\chi}$ (d) $y = (0.05)^{\chi}$
- (9) In \triangle ABC, if a = b = 8 cm. and the perimeter of \triangle ABC = 26 cm. , then m (\angle C) \approx
 - (a) 35.3°

- (b) 52.3°
- (c) 77.4°
- (d) 108°
- (10) In \triangle ABC, if m (\angle A) = 30° and a = 6 cm., then $\frac{b}{\sin B}$ =cm.
 - (a) 3

- (b) 6
- (c) $\frac{1}{5}$
- (d) 12

- (11) $\lim_{x \to 1} \frac{x^5 1}{x 1} = \dots$

- (c) 4
- (d) 20

- (12) In any triangle LMN, $\frac{\ell}{\sin L} = \dots$

- (b) $\frac{n}{\sin M}$
- (c) $\frac{m+n}{\sin N + \sin N}$ (d) 3 r

- (13) $\lim_{x \to \infty} \frac{\sqrt{4x^2 + 1}}{x 2} = \dots$

- (c) $\frac{5}{2}$
- (d) 2

- (14) $\lim_{x \to 0} (2x^2 + 3) = \cdots$
- (a) 2

- (d)7
- (15) In \triangle ABC, if $2 \sin A = 3 \sin B = 4 \sin C$, then $a : b : c = \dots$
 - (a) 2:3:4
- (b) 4:3:2
- (c) 3:4:6
- (d) 6:4:3
- (16) In \triangle ABC, if $4 \sin A = 3 \sin B = 6 \sin C$, then m (\angle C) \simeq
 - (a) 89°

- (b) 29°
- (c) 57°
- (d) 82°

(17) The solution set in I	\mathbb{R} of the equation : 2^{2X}	$-12 \times 2^{X} + 2^{5} = 0$	equals	
(a) $\{4, 8\}$	(b) $\{2, 3\}$	(c) $\{16, 2\}$	(d) $\{1, 4\}$	
(18) The function $f:f$	$(X) = a^{X}$ is increasing if			
(a) $a > 0$	(b) $a > 1$	(c) $a = 1$	(d) $0 < a < 1$	
(19) ABC is an equilatera	al triangle its side lengt	$h = 5\sqrt{3}$ cm., then t	he diameter length of its	
circumcircle equals	cm.			
(a) $5\sqrt{3}$	(b) $10\sqrt{3}$	(c) 10	(d) 5	
(20) $\log_{5} 49 \times \log_{8} 5 \times 1$	$\log_9 8 \times \log_7 9 = \dots$	******		
(a) log 100	(b) log 7	(c) log 5	(d) log 2	
(21) If $f: \mathbb{R} \longrightarrow \mathbb{R}$, w	here $f(X) = (a+1)X +$	b-2 and $f(X)$ map	s each real number to	
itself, then $(a, b) =$				
(a) (0,3)	(b) $(0, -3)$	(c)(0,2)	(d) $(-1, 2)$	
(22) The type of the func	tion $f: f(x) = \frac{\sin x}{x}$ is			
(a) even.	X	(b) odd.		
(c) neither odd nor even.		(d) both odd and even.		
(23) $\lim_{X \to \frac{\pi}{4}} \frac{\tan X}{X} = \cdots$				
(a) $\frac{\pi}{4}$	(b) I	$(c)\frac{4}{\pi}$	(d) does not exist.	
(24) $\lim_{x \to 1} \frac{2x+a}{x+1} = 5$,	then a =	***		
(a) 2	(b) 5	(c) 8	(d) 10	
(25) In any triangle XYZ	$\mathcal{X}^2 + \mathcal{Y}^2 - 2 \mathcal{X} \mathcal{Y} \cos z$	Z =		
(a) X ²	(b) y ²	(c) z ²	(d) z	
(26) If $f(x) = \frac{\sqrt{x^2 - 2x}}{x - 1}$	$\frac{+1}{}$, then the range of the	he function f is		
(a) {1}		(c) [-1,1[(d) $\{-1, 1\}$	
(27) $\lim_{x \to 1} \frac{x^2 - 1}{x - 1} = \dots$				
(a) 0	(b) 1	(c) 2	(d) 3	
(28) In ∆ ABC , m (∠ A)	$: m (\angle B) : m (\angle C) =$	$3:5:4$, then $c^2:a$	² =	
$(a)\sqrt{6}:2$	(b) 2:3	(c) 4:3	(d) 3:2	

Second **Essay questions**

Answer the following questions:

- Graph the curve of the function f where f(x) = |x-3|, deduce the range and monotony of the function and tell whether it is even , odd or otherwise.
- Find the solution set of the following equation in \mathbb{R} : $\log_2 x + \log_2 (x + 1) = 1$
- Find the value of the following: $\lim_{x \to \infty} \frac{4-3x^2}{\sqrt{x^4+5}}$
- Find the value of the following: $\lim_{x \to -1} \frac{2x^3 x^2 2x + 1}{x^3 + 1}$

El-Menia Governorate



Minia Governmental Language School

Multiple choice questions First

Choose the correct answer from the given ones:

- (1) The range of the function f: f(x) = |x| is
 - (a) [0,∞[
- (b)]0,∞[
- $(c) \infty, 0$
- $(d) \infty, 0$
- (2) The curve of the even function is symmetric about the straight line
 - (a) y = x

- (d) v = -x
- (3) The S.S. of the inequality $|3-2x| \le 1$ in \mathbb{R} is
 - (a) [1,2]

- (b)]1,2[
- (c) $\mathbb{R} [1, 2]$ (d) $\mathbb{R} [1, 2]$
- (4) The range of the function $f: f(X) = \frac{15}{x} + 2$ is
 - (a) T

- (b) $\mathbb{R} \{2\}$
- (c) {2}
- $(d) \mathbb{R} \{0\}$
- (5) The S.S. of the equation $|2 \times -1| = 5$ in \mathbb{R} is
 - (a) $\{3\}$

- (b) $\{-2\}$
- (c) Ø
- (d) $\{3, -2\}$
- (6) The point of symmetry of the curve of the function $f: f(X) = X^3$ is
 - (a) (1, 1)

- (b)(0,0)
- (d) (0,1)
- (7) The domain of the function $f: f(X) = \frac{2X}{X^2 4}$ is
- (a) $\mathbb{R} \{-2, 2\}$
- (b) $\mathbb{R} \{-2, 0, 2\}$ (c) \mathbb{R}
- (d) $\mathbb{R} \{4\}$
- (8) The function $f: f(x) = a^x$ is decreasing if
 - (a) a = 1

- (b) a > 1
- (c) 0 < a < 1
- (d) a = -1

(9) If $3^{X+1} - 3^X = 54$, then $X = \cdots$		
(a) 1	(b) 2	(c) 3	(d) 4
(10) If $f(x) = 3^{x+2}$, t	hen $f(X+1) \times f(-X) =$	=	
(a) 27	(b) 81	(c) 243	(d) 729
(11) If $\log 3 = X \cdot \log 5$	$= y$, then $log 15 = \cdots$		
(a) $X + y$	(b) $X - y$	(c) X y	(d) $\frac{x}{y}$
(12) If $3^{X-2} = 2^{X-2}$,	then $X = \cdots$,
(a) 3	(b) - 2	(c) 0	(d) 2
(13) The solution set of	the equation : $\chi \frac{4}{3} = 81$ in	n R is	
(a) $\{-27, 27\}$	(b) $\{9, -9\}$	(c) {9}	(d) $\{27\}$
(14) If $2^{X-3} = 1$, then	<i>X</i> = ······		
(a) - 3	(b) 3	(c) 1	(d) zero
(15) $\lim_{x \to 1} \frac{2x-4}{x-2} = \cdots$			
(a) 1	(b) 2	(c) - 2	(d) zero
(16) $\lim_{x \to 4} \frac{x^2 + 7x + 1}{x^2 - 6x + 5}$	$\frac{5}{3} = \frac{15}{2}$, then b =	****	
(a) - 44	(b) 7	(c) - 8	(d) 8
(17) $\lim_{y \to 2} \frac{y^5 - 32}{y - 2} = \cdots$			
(a) 31 y^4	(b) 32×2^4	(c) 64	(d) 5×2^4
(18) $\lim_{y \to 0} \frac{x^2 + x}{x} = \cdots$			
(a) zero	(b) 1	(c) 2	(d) 3
(19) $\lim_{x \to \infty} \left(\frac{3 x^2 + 2 x}{x^2 - 3 x} \right)$	$\left(\frac{1}{1} + \frac{1}{2}\right)^4 = \cdots$		
(a) 3	(b) 9	(c) 27	(d) 81
$(20) \lim_{X \to \infty} \frac{k X}{3 X + 1} = 4$, then $k = \cdots$		
(a) 16	(b) 12		(d) $\frac{4}{3}$
(21) In \triangle XYZ, $\frac{x}{\sin X}$ =	6, then the length of dian	neter of its circumcirc	ele = ····· length unit
(a) 4	(b) 12		(d) 9
(22) In \triangle ABC : if 2 sin	$A = 3 \sin B = 4 \sin c$, th	en a : b : c =	
(a) 2:3:4	(b) 4:3:2	(c) 3:4:6	(d) 6:4:3
	greatest angle of the trian	ngle whose side leng	ths are 3 cm., 5 cm. and
7 cm. is			
(a) 150°	(b) 120°	(c) 60°	(d) 30°

- - (a) 24.72

- (b) 26.3
- (c) 28.88
- (d) 30
- (25) In \triangle ABC, if $4 \sin A = 3 \sin B = 6 \sin C$, then m (\angle C) = (to nearest degree)
 - (a) 89°

- (b) 29°
- (c) 57°
- (d) 82°
- (26) If ABC is a triangle in which a = 4 cm., $b = 4\sqrt{3} \text{ cm.}$, c = 8 cm., then sine of its smallest angle =
 - (a) $\frac{1}{2}$

- (b) $\frac{\sqrt{3}}{2}$
- (c) 1
- (d) zero
- - (a) 14

- (b) $14\sqrt{3}$
- (c) 8
- (d) 28 \(\sqrt{3}\)

- (28) In any $\triangle XYZ : X^2 + y^2 2 X y \cos z = \dots$
 - (a) χ^2

- (b) y^2
- (c) z^2
- (d) z

Second Essay questions

Answer the following questions:

- 1 Find in \mathbb{R} the solution set of : $|X-3| \le 4$
- **2** Find the domain of $f: f(X) = \log_4 (4 X)$
- 3 Find: $\lim_{x \to 2} \frac{x^3 8}{x^2 5x + 6}$
- 4 Find: $\lim_{x \to \infty} \frac{2x-9}{|3x|+7}$

10 Aswan Governorate



Aswan Educational Administration M.M. Yaqoub Language School

First Multiple choice questions

Choose the correct answer from the given ones:

- (1) Vertex of function $f: f(x) = (x-4)^2 + 2$ is
 - (a) (-4, 2)
- (b) (2,4)
- (c) (4, -2)
- (d)(4,2)

- (2) $\lim_{x \to -2} (3x^2 + x 4)$ is
 - (a) 3

- (b) 12
- (c) 9
- (d) 6

- (3) If $3^{X-3} = 4^{X-3}$, then $X = \dots$
 - (a) $\{9\}$

- (b) $\{-3\}$
- (c) {zero}
- $(d) \{3\}$

- (4) Domain of $f: f(x) = \frac{x+2}{x^2-4}$ is
 - (a) $\mathbb{R} \{2\}$
- (b) $\mathbb{R} \{-2\}$ (c) $\mathbb{R} \{2, -2\}$ (d) \emptyset
- (5) In \triangle ABC which is drawn in a circle, then $\frac{1}{2 r} = \cdots$

- $\frac{b}{\sin B} \qquad \qquad (c) \frac{c}{\sin C}$

- (6) $\lim_{x \to 0} \frac{x^2 x}{x}$ is
 - (a) zero

- (c) ∞
- (d) 1
- (7) In \triangle ABC, if a = 5, b = 7, c = 8, then $m (\angle B) \simeq \dots$
 - (a) 90°

- (b) 80°
- (c) 70°
- (d) 60°
- (8) Diameter length of circumcircle of triangle ABC in which m (\angle A) = 60° $, a = \sqrt{3} \text{ cm. is } \dots \text{ cm.}$
- (b) $2\sqrt{3}$ (c) 2
- (d) 1/3

- (9) $\lim_{x \to 2} \sqrt{3x+3} = \cdots$

- (b) 3
- (c) 3
- $(d) \pm 3$

- (10) Solution set of: |X| + 3 = 0, is
 - $(a) \pm 3$

- (b) 3
- (d) Ø

- (11) Type of function $f: f(x) = 2 x^2$ is
 - (a) even.

(b) odd.

(c) neither even nor odd.

- (d) increasing.
- (12) Monotony of function $f: f(X) = \left(\frac{1}{5}\right)^X$, is
 - (a) increasing.

(b) decreasing.

(c) increasing and decreasing.

(d) constant.


- (13) $\lim_{x \to a} \frac{x^n a^n}{x a}$ is
 - (a) an a-1

- (b) na^{1-n}
- (c) na 1 a
- (d) na n-1

(14) In the opposite figure:

Rule of the function is

- (a) f(x) = |x-1| + 3
 - (b) f(x) = 3 |x + 1|
 - (c) f(x) = 3 |x 1|
 - (d) f(x) = |1 x| + 3



(15) In ∆ ABC if m (∠ A)	= 30° and a = 6 cm.	then $\frac{b}{\sin B} = \dots$	•
(a) 11	(b) 21	(c) 13	(d) 12
(16) $\lim_{X \to 1} \frac{X^2 + X - 2}{X - 1} = \cdots$			
(a) 1	(b) 2	(c) 3	(d) 4
(17) In \triangle ABC: $b^2 + c^2 -$	a ² =		
(a) 2 bc cos A	(b) 2 ac cos B	(c) 2 bc cos C	(d) 2 ab cos C
(18) The range of the func	tion $f: f(x) = -(x)^2$	is	
(a) $]-\infty$, 0[(b) $]0, \infty[$	(c) [0,∞[(d) $]-\infty,0]$
(19) If: $\log_2(x) + \log_2(x)$	$(X+1) = 1$, where $X \in$	$\exists \mathbb{R}$, then $X = \cdots$	
(a) $\{-1, 2\}$	(b) {1}	(c) {2}	(d) $\{-2, 1\}$
(20) $\lim_{x \to 0} \frac{(x+1)^{17}-1}{x} =$	***********		
(a) 15		(c) 18	(d) 17
(21) Symetric point of the	curve $f : f(x) = 3 - \frac{1}{2}$	$\frac{1}{2-X}$	
(a) $(3, -2)$	(b) $(-2,3)$	(c) $(-2, -3)$	(d) (2,3)
(22) $\log_5 \sqrt{5} = \dots$			
(a) 2	(b) 5	(c) $\frac{1}{2}$	(d) - 1
(23) In Δ ABC if a : b : c =	$= 3:2:2$, then $\cos A$	=	
(a) $\frac{1}{2}$	1.	(c) $\frac{3}{4}$	$(d) - \frac{1}{8}$
(24) $\lim_{x \to 2} \frac{x^5 - 32}{x - 2} = \dots$			
(a) 64	(b) 46	(c) 80	(d) 82
(25) If $f: f(x) = \frac{1}{3}$, the	$\inf\left(\frac{1}{3}\right) = \dots$		
(a) 1 (26) $\lim_{x \to \infty} \frac{\sqrt{4x^2 - 1}}{x - 2} = 0$	(b) $\frac{1}{9}$	(c) 3	(d) $\frac{1}{3}$
(a) 4	(b) 5	(c) 1	(d) 2
(27) Range of the function	$f: f(X) = X \text{ is } \cdots$	*****	
(a) $\mathbb{R}-\{1\}$		(c) IR	
(28) The two curves of the	e two functions $f:f(0)$	$(x) = 2^{x}$ and $g : g(x)$	$=\left(\frac{1}{2}\right)^X$ intersect
at $X = \cdots y =$			
(a) $(-2,0)$	(b) $(0, -2)$	(c)(0,1)	(d) (1,0)

Second Essay questions

Answer the following questions:

- 1 Find: $\lim_{x \to \infty} \left(5 \frac{5}{x^3} \right)$
- Use the curve of $f: f(X) = X^3$ to graph $g: g(X) = X^3 3$ From the graph deduce domain and its range.
- Use the curve of the function f where $f(X) = \frac{1}{X-1}$ to represent g where g(X) = f(X) + 2Find:
 - (a) Monotony of the function g
 - (b) Range of g
- Find the perimeter of \triangle ABC in which a = 8 cm. , b = 6 cm. and $m (\angle C) = 48^{\circ}$

Model

Interactive test



First

Multiple choice questions


Choose the correct answer from the given ones:

11 The range of the given function in the opposite figure is



(b)
$$\{1, -1\}$$

(c)
$$\{-1\}$$



2 If $5^{X-3} = 4^{3-X}$, then $X = \dots$

(a)
$$\frac{5}{4}$$

(c)
$$\frac{4}{5}$$

 $\lim_{x \to \infty} \frac{2x+3}{5x^2+4}$



(b) zero




- (d) $\frac{2}{5}$
- In \triangle ABC, if $4 \sin A = 3 \sin B = 6 \sin C$, then m (\angle C) \simeq
 - (a) 89°


(b) 29°

- (c) 57°
- (d) 82°
- - (a) 7


(b) 9

- (c) 13
- **6** If $f: \mathbb{R} \longrightarrow \mathbb{R}$ where $f(x) = x^3$, then the figure which represents the function fis









- The solution set in \mathbb{R} of the equation: $2^{2^{X}} 12 \times 2^{X} + 2^{5} = 0$ equals
 - (a) $\{4, 8\}$

(a)

(b) $\{2,3\}$

(b)

- (c) $\{16, 2\}$
- $(d) \{1,4\}$

- $\lim_{X \to 0} \frac{(X+2)^5 32}{X} = \dots$
 - (a) 25

(b) 64


- (c) 80
- (d) 100

- The opposite figure represents the curve of the function f, then $\lim_{x \to 2} f(x) = \cdots$
 - (a) 1

(b) - 1

(c) 2

(d) does not exist.



- 10 The function $f: f(X) = a^X$ is increasing if
 - (a) a > 0

- (b) a > 1
- (c) a = 1
- (d) 0 < a < 1

- If $X = 5 + 2\sqrt{6}$, then $\log \left(X + \frac{1}{X}\right) = \dots$
 - (a) 1

- (b) $5 2\sqrt{6}$
- (c) 10
- (d) $5 + 2\sqrt{6}$
- ABC is an equilateral triangle, its side length = $5\sqrt{3}$ cm., then the diameter length of its circumcircle equals cm.
 - (a) $5\sqrt{3}$

- (b) $10\sqrt{3}$
- (c) 10
- (d) 5

- $\log_5 49 \times \log_8 5 \times \log_9 8 \times \log_7 9 = \dots$
 - (a) log 100

- (b) log 7
- (c) log 5
- (d) log 2

- $\lim_{x \to 1} \frac{x^7 1}{x 1} = \dots$
 - (a) 35

(b) 7

- (c) 42
- (d) 1
- **15** The solution set of the equation : $\log_3 x \times \log_2 3 = 5$ in \mathbb{R} is
 - (a) $\{32\}$

(b) {5}

- (c) $\{3\}$
- (d) $\{2\}$
- In \triangle ABC, $m(\angle A) : m(\angle B) : m(\angle C) = 3 : 5 : 4$, then $c^2 : a^2 = \dots$
 - (a) $\sqrt{6}:2$


(b) 2:3

- (c) 4:3
- (d) 3:2

11 In the opposite figure :

The solution set of the inequality : f(X) < g(X) in \mathbb{R} is

- (a) $\{-4,0\}$
- (b) [-4,0]
- (c) $\mathbb{R} [-4, 0]$
- (d)]-4,0[



- 18 The type of the function $f: f(X) = \frac{\sin X}{x}$ is
 - (a) even.

(b) odd.

(c) neither odd nor even.

(d) both odd and even.

- $\lim_{X \to \frac{\pi}{4}} \frac{\tan X}{X} = \dots$
 - (a) $\frac{\pi}{4}$

- (b) 1
- $(c)\frac{4}{\pi}$
- (d) does not exist.

- 20 If $x^{\frac{3}{2}} = 8$, then $x = \dots$
 - (a) 2

- (b) 4
- (c) 8
- (d) 9

- $\lim_{x \to 2} \frac{x^3 7x + 6}{3x^2 8x + 4} = \dots$
 - (a) $\frac{4}{5}$

- (b) $\frac{2}{3}$
- (c) $\frac{5}{4}$
- (d) $\frac{3}{2}$

- 22 If $\lim_{x \to 1} \frac{2x + a}{x + 1} = 5$, then $a = \dots$
 - (a) 2

- (b) 5
- (c) 8
- (d) 10
- In any triangle XYZ, $x^2 + y^2 2x$ y cos Z =
 - (a) x^2

- $(c) z^2$
- (d) Z
- The number of possible solutions for the triangle ABC where : $m (\angle A) = 60^{\circ}$, b = 3 cm. , a = 5 cm. is
 - (a) 1

(b) 2

(c) zero

- (d) infinite number.
- 25 If $(\frac{1}{2})^{a^2-a-2} = 1$ where a > 0, then $a = \dots$
 - (a) 1

- (b) 3
- (c) 2
- (d)3
- If $f(x) = \frac{\sqrt{x^2 2x + 1}}{x 1}$, then the range of the function f is
 - (a) $\{1\}$

- (b) IR
- (c) [-1,1[(d) $\{-1,1\}$

In the opposite figure:


The length of AB ≈ ····· cm.

(a) 6

(b) 7

(c) 8

(d) 9




- In \triangle XYZ, the expression $\frac{\chi^2 + y^2 z^2}{2 \chi y}$ equals
 - (a) cos X
- (b) cos Y
- (c) cos Z
- (d) sin Z

Second Essay questions

Answer the following questions:

- Use the curve of the function f where $f(X) = \frac{1}{X}$ to represent the function g: g(X) = f(X-2) + 2 and from the graph determine the range and discuss its monotony.
- 2 Find: $\lim_{x \to \infty} \frac{\sqrt{4x^2 + 1}}{5x + 2}$
- Find the solution set in \mathbb{R} of the inequality : $\sqrt{4x^2 12x + 9} \le 9$
- From the opposite figure , find :
 - (1) $\lim_{X \to -2} f(X)$
 - (2) f(-2)
 - (3) $\lim_{X \to 0} f(X)$
 - (4) f(0)



Model

2

Interactive test 2



First Multiple choice questions

Choose the correct from the given ones:

- 11 The range of the function f: f(X) = |X| is
 - (a) $[0, \infty[$
- (b) $]0, \infty[$
- (c) $]-\infty,0]$
- (d) $]-\infty,0[$

- $\lim_{X \to \infty} \left(\frac{3}{5}\right)^{\frac{1}{X}} = \cdots$
 - (a) 1

- (b) 1
- (c) $\frac{3}{5}$
- (d) ∞

- 3 $\lim_{x \to 1} \frac{x^5 1}{x 1} = \dots$
 - (a) 5

- (b) 1
- (c) 4
- (d) 20

- - (a) 2

(b) 3

- (d) 6
- 5 If f is an odd function and $X f(X) + X^3 f(-X) = 2$, then $f(2) = \cdots$
 - (a) 3

(b) $\frac{1}{2}$

- $(c) \frac{1}{3}$
- (d) 3

- **6** In \triangle XYZ, $\frac{X^2 + y^2 z^2}{2 X y} = \dots$
 - (a) cos X

- (b) cos Y
- (c) cos Z
- (d) sin Z

- If $f(x) = 3^x$, then the solution set in \mathbb{R} of the equation $f(2 X) - 28 f(X) + f(3) = \text{zero equals} \dots$
 - (a) $\{1, 27\}$
- (b) $\{1,3\}$
- (c) $\{0,3\}$ (d) $\{3\}$
- The logarithmic form that equivalent to the exponential form: $2^7 = 128$ is
 - (a) $\log_2 128 = 7$

(b) $\log_2 7 = 128$

(c) $\log_7 128 = 2$

- (d) $\log_7 2 = 128$
- The curve of the even function is symmetric about the straight line
 - (a) y = X

 $(b) \overline{yy}$

- (c) XX
- (d) y = -X
- In \triangle LMN, $\frac{\sin L}{3} = \frac{2 \sin M}{3} = \frac{\sin N}{4}$, then $\ell : m : n = \dots$
 - (a) 6:8:3
- (b) 3:6:8
- (d) 6:3:8
- In \triangle ABC, c = 7 cm., $m (\angle A) = 70^{\circ}$, $m (\angle B) = 40^{\circ}$, then $b \simeq \cdots$ cm.
 - (a) 3.7

(b) 4.8

- (c) 8.4
- (d) 7.3

- 12 If $\lim_{x \to a} \frac{a x}{3} = 12$, then $a = \dots$
 - $(a) \pm 12$

 $(b) \pm 6$

(c) 3

- (d) 3
- 13 The range of the function $f: f(x) = \frac{x-2}{2-x}$ equals
 - (a) IR

- (b) ℝ {2}
- (c) $\mathbb{R} \{-2\}$
- (d) $\{-1\}$

- 14 If $\log 3 = x$, $\log 7 = y$, then $\log 21 = \dots$
 - (a) X y

- (b) X + y
- (c) X y
- $(d)\frac{x}{y}$

- 15 $\log_3 5 \times \log_2 3 \times \log_5 16 = \dots$
 - (a) 30

- (b) 15
- (c) log 10000
- (d) log₃₀ 240
- The curve of the function $g: g(X) = X^2 + 4$ is the same as the curve of $f: f(X) = X^2$ by translation 4 units in the direction of
 - (a) \overrightarrow{Ox}

- (b) \overrightarrow{Ox}
- (c) \overrightarrow{Oy}
- (d) Oy


- $\lim_{x \to 5} \frac{\sqrt{x-1}-2}{x-5} = \dots$
 - (a) $\frac{4}{3}$

- (b) $\frac{3}{4}$
- (c) 4
- (d) $\frac{1}{4}$
- 18 The function f where $f(X) = a^X$ is decreasing on its domain if
 - (a) a = 1

- (b) a > 1
- (c) 0 < a < 1
- (d) a = -1

- $\lim_{X \to 0} \frac{(4X+1)^9 1}{3X} = \dots$
 - (a) $\frac{3}{4}$

- (b) $\frac{4}{3}$
- (c) 9
- (d) 12
- The solution set in \mathbb{R} of the equation : |x-7| = 2 is
 - (a) $\{9,5\}$
- (b) $\{7,3\}$
- (c) Ø
- (d) $\{3, -3\}$



- If the perimeter of \triangle ABC = 33 cm. , $\sin A + \sin C = \frac{2}{3}$, $\sin B = \frac{1}{4}$, then AC = cm.
 - (a) 6

- (b) 9
- (c) 12
- (d) 15

- - (a) 28.3

- (b) 38.3
- (c) 3.8

(d) 28.4

- $\lim_{x \to \infty} (5 + 3 x^2 + x) = \dots$
 - (a) not exist.
- (b) 5
- (C) 00

- (d)9
- 25 ABCD is a parallelogram, $m (\angle A) = 50^{\circ}$, $m (\angle DBC) = 70^{\circ}$, BD = 8 cm. , then the perimeter of the parallelogram ABCD to the nearest cm. = cm.
 - (a) 38

- (b) 30
- (c) 19

(d) 48

26 The solution set of the inequality $|X-1| \le 3$ is

(a)
$$[-2,4]$$

(b)
$$]-2,4[$$

(b)
$$]-2,4[$$
 (c) $]-2,4[$

(d) $\mathbb{R} - [-2, 4]$

- In \triangle ABC, $\cos(A + B) = \cdots$
 - (a) cos C

- (b) cos C
- (c) sin C
- $(d) \sin C$

28 In the opposite figure:

M is the centre of the circle


 $, BM = 6 \text{ cm.}, \text{ then } AB = \dots \text{ cm.}$

(a) 6 sin 50°

(b) 12 sin 50°

(c) 6 cos 50°

(d) 12 cos 50°



Essay questions Second

Answer the following questions :

- 11 If $x = 5 + 2\sqrt{6}$, find in the simplest form the value of $\log(\frac{1}{x} + x)$ without using calculator.
- 2 Use the curve of the function $f: f(x) = \frac{1}{x}$ to graph the curve of the function g: g(X) = $\frac{1}{X-2}$ + 3, from the graph state the domain and range of g and the monotony and its type whether it is even , odd or otherwise.
- 3 Find: $\lim_{x \to 1} \frac{(x+2)^4 81}{x-1}$
- $\lim_{x \to \infty} \frac{6x 4x^3}{2 7x^3}$

Model

Interactive test 3



First Multiple choice questions

Choose the correct answer from the given ones:

- If $f(X) = 7^{X+1}$, then the solution set of the equation : f(2X-1) + f(X-2) = 50in R equals
 - (a) $\{1\}$

- (b) $\{1, -1\}$ (c) $\{1, -50\}$ (d) $\{7, -50\}$

- If $\log 3 = X$, $\log 5 = y$, then $\log 15 = \cdots$
 - (a) χ y
- (c) X + y
- (d) X y

- $\lim_{x \to \infty} \frac{5 + x^{-2}}{1 + 3 x^{-2}} = \dots$

- (c) $\frac{5}{3}$
- (d)5

- 4 If $f(x) = 5^x$, then $f(-2) = \cdots$
 - (a) 2

- (b) 5
- (c) $\frac{1}{25}$
- (d) $\frac{1}{5}$
- The domain of the function $f: f(X) = \log_3 (X-2)$ is $X > \dots$
 - (a) 3

- (b) 5
- (c) 1

(d)2

- $\log 25 + \frac{\log 8 \times \log 16}{\log 64} = \dots$
 - (a) log₂ 16
- (b) log₅ 25
- (c) log 4
- (d) log 10


In the opposite figure:

If the perimeter of \triangle ABC = 42 cm.,

the circle touches the sides of the triangle

internally, then: $m (\angle B) = \cdots$

- (a) 53° 8
- (b) 67° 23
- (c) 36° 53
- (d) 32° 37



(a) zero

- (b) 1
- (c) 2

(d) ∞

- $\lim_{x \to \infty} \frac{\sqrt{x^2}}{x} = \dots$
 - (a) zero

- (b) 2
- (C) 00
- (d) 1

- $\lim_{x \to 5} \frac{x^2 8x + 15}{x^2 10x + 25} = \dots$
 - (a) does not exist.
- (b) zero
- (c) 2
- (d) 3
- 11 The included area between the curves of the two functions f: f(x) = |x+3| 2 $g: g(X) = \text{zero is } \dots \text{square units.}$
 - (a) 2

- (b) 3
- (c) 4
- (d) 5

- 12 If $\log_3 y = X$, then the exponential form is
 - (a) $y = x^3$
- (b) $X = y^3$ (c) $X = 3^y$
- (d) $y = 3^{X}$
- 13 If f is an odd function on [-X, X], then $f(-X) + f(X) = \cdots$
 - (a) 2 X

- (b) undefined.
- (c) -2x
- (d) zero
- In \triangle ABC, if $2 \sin A = 3 \sin B = 4 \sin C$, then $a : b : c = \dots$
 - (a) 2:3:4
- (b) 4:3:2
- (c) 3:4:6
- (d) 6:4:3

- 15 $\lim_{x \to 2} \frac{x^5 32}{x^2 + 3x 10} = \dots$
 - (a) $\frac{16}{7}$

- (b) $\frac{80}{7}$
- (c) $\frac{7}{80}$
- (d) $\frac{7}{16}$
- 16 The radius length of the circumcircle of the triangle ABC in which m (\angle A) = 30° , a = 10 cm. equals
 - (a) 10 cm.
- (b) 20 cm.
- (c) 5 cm.
- (d) 40 cm.
- - (a)]0,2[
- (b)]-∞,0[
- (c) $\mathbb{R} [0, 2]$
- (d)]0,∞[
- **1B** The solution set of the equation : $\log_{\chi} 81 = 4$ in \mathbb{R} is
 - (a) $\{-3\}$
- (b) $\{3\}$ (c) $\{3, -3\}$ (d) $\{9\}$
- 19 The solution set of the equation : |x + 2| = -2 in \mathbb{R} is
 - (a) Ø

- (b) R
- (c) $]-\infty, -2[$ (d) $]-\infty, -2[$

- The measure of the greatest angle in the triangle whose side lengths are 3 cm., 5 cm.
 - , 7 cm. equals
 - (a) 150°

- (b) 120°
- (c) 60°
- (d) 30°

211 In the opposite figure:


If
$$CD = CB = 6 \text{ cm}$$
.


- then $\tan \theta = \cdots$
- (a) $\frac{3}{4}$

(b) $\frac{4}{3}$

(c) $\frac{1}{2}$

(d)2






23 The rule of the function shown

in the opposite figure is $f(X) = \cdots$

- (a) $(X-2)^2 + 1$ (b) $-(X-2)^2 + 1$
- (c) $-(x-1)^2 + 2$ (d) $(-x+1)^2 + 1$



- 24 In \triangle ABC, $a^2 + b^2 c^2 = \dots$
 - (a) cos A


- (b) a b cos C
- (c) cos C
- (d) 2 a b cos C
- **25** The solution set of the equation : $\chi^{\frac{2}{3}} = 25$ in \mathbb{R} is
 - (a) {5}

- (b) $\{5, -5\}$
- (c) {125}
- (d) $\{125, -125\}$

In the opposite figure:

$$\lim_{x \to 2} f(x) = \cdots$$

- (a) zero
- (b) not exist.
- (c) 2
- (d) 1



The number of possible solutions of \triangle ABC in which a = 8 cm., b = 10 cm.

, m (
$$\angle$$
 A) = 42° is

(a) 1

- (b) 2
- (c) infinite number. (d) zero.


28 In the opposite figure :

(a) 6

(b) 7

(c) 8

(d) 9



Essay questions Second

Answer the following questions:

1 Prove that:
$$\frac{2^{x} \times 9^{x+1}}{3 \times (18)^{x}} = 3$$

2 Graph the function
$$f: f(x) = \begin{cases} |x|, & x \le 0 \\ x^3, & x > 0 \end{cases}$$
, from the graph state the range of the function and discuss its monotony.

3 Find:
$$\lim_{x \to \infty} \frac{(x+1)(5x-3)}{x^2+3}$$

$$\lim_{x \to 2} \frac{5 \, x - 10}{4 \, x - 8}$$

Model 4

Interactive test 4



First Multiple choice questions

Choose the correct answer from the given ones:

- $\lim_{x \to 0} \frac{x^7 1}{x + 1} = \dots$
 - (a) 2

- (b) 5
- (c) 1

(d) - 1

- 2 In \triangle ABC, $\frac{b^2 + c^2 a^2}{2bc} = \cdots$
 - (a) cos A

- (b) cos B
- (c) cos C
- (d) sin A
- The solution set in \mathbb{R} of the inequality : $|x-1| \ge 3$ equals
 - (a) $\mathbb{R}]-2,4[$ (b) [-2,4] (c) $\mathbb{R} [-2,4]$ (d)]-2,4[

- 4 $\lim_{X \to -1} \frac{x^2 + x}{x^3 + 1} = \dots$
 - (a) zero

- (b) $-\frac{1}{3}$
- (c) 1
- (d) does not exist.
- The radius length of the circumcircle of \triangle XYZ in which $\mathcal{X} = 20 \sin X \text{ cm}$. equals cm.
 - (a) 5

- (b) 10
- (c) 20
- (d) 40
- 6 Which of the following functions represents an increasing exponential function on its domain R?
 - (a) $y = 3 (1.05)^{x}$
- (b) $y = 3 \left(\frac{1}{1.05}\right)^{x}$ (c) $y = 3 + (0.5)^{x}$ (d) $y = (0.05)^{x}$
- The solution set of the equation : $\log 5 X = -1$ in \mathbb{R} is
 - (a) $\left\{ \frac{1}{10} \right\}$
- (b) $\left\{ \frac{1}{50} \right\}$
- (c) {1}
- $(d) \{50\}$
- The measure of the smallest angle in \triangle ABC in which, a = 8 cm., b = 7 cm., and its perimeter is 21 cm, approximately equals
 - (a) 32° 34
- (b) 42° 34
- (c) 36° 34
- (d) 46° 34

- If $5^{x} = 17$, then the value of x to the nearest two decimals equals
 - (a) 1.34

- (b) 1.32
- (c) 1.76
- (d) 1.67

- $\lim_{h \to 0} \frac{(2-3h)^7 128}{4h} = \dots$
 - (a) 336

- (b) 336
- (c) 623
- (d) 633
- If the curve of the function $f: f(X) = \log_4 (1 aX)$ passes through the point $(\frac{1}{8}, \frac{-1}{2})$ • then a =
 - (a) 3

- (b) 2
- (c) 4

- (d) 8
- 12 The solution set of the equation : $\chi^{\frac{4}{3}} = 81$ in \mathbb{R} is
 - (a) $\{27, -27\}$
- (b) $\{9, -9\}$ (c) $\{9\}$
- (d) $\{27\}$

- 13 $\lim_{x \to \infty} \frac{(12)^{\frac{1}{X}}}{x+7} = \dots$
 - (a) $\frac{12}{7}$

- (b) ∞
- (c) 1

(d) zero

14 In the opposite figure :


If BC = 10 cm., then the perimeter of \triangle BDC \simeq cm.

(a) 60

(b) 62

(c) 64

(d) 67



- 15 If $3^{X-2} = 2^{X-2}$, then $X = \dots$
 - (a) 3


- (b) 2
- (c) zero
- (d) 2
- 16 The domain of the function $f: f(X) = \frac{1}{|X|-3}$ is
 - (a) $\{3, -3\}$

- (b) [-3,3] (c) $\mathbb{R} [-3,3]$ (d) $\mathbb{R} \{-3,3\}$
- 11) The vertex of the curve of the function $f: f(x) = (2-x)^2 + 3$ is
 - (a)(2,3)
- (b) (2, -3) (c) (-2, 3)
- (d) (-2, -3)

- $\lim_{X \to 0} \lim_{x \to 0} \frac{(2 X + 1)^2 1}{x} = \dots$
 - (a) 4

- (b) 3
- (c) 4
- (d) 2

19 The curve represented in the opposite figure is symmetric about the straight line whose equation is



(a) X = 0

- (b) y = 0
- (c) y = -2
- (d) X = 2
- If \angle A supplements \angle C, then $\cos A + \cos C = \cdots$
 - (a) 1

- (b) zero
- (c) $\frac{1}{2}$
- (d) 1

- $\lim_{x \to 0} \frac{5+2x}{\cos 3x} = \cdots$
 - (a) 5


- (b) 3
- (c) 2
- (d) $\frac{5}{3}$
- If $\log_2 x = 4$, then the exponential form that equivalent to it is
 - (a) $2^{\tilde{x}} = 4$
- (b) $X = 2^4$
- (c) $X^2 = 4$
- (d) $4^{x} = 2$


- 23 If $\frac{a+b}{13} = \frac{b+c}{11} = \frac{c+a}{12}$, then $\cos A = \dots$
 - (a) $\frac{1}{5}$


- (b) $\frac{5}{7}$ (c) $\frac{19}{35}$
- (d) $\frac{4}{11}$
- The solution set of the equation: $(\log_2 x)^2 2 \log_2 x = 3$ in \mathbb{R} is
 - (a) {16}


- (b) $\{8\}$
- (c) {8,0.5}
- (d) $\{16, 0.5\}$
- If g is a real function whose domain is [-2, 3], then the domain of n : n (x) = g(x-2)is
 - (a) [-2,3]
- (b) [-4,1]
- (c) [0,5]
- (d) IR
- **26** If the radius length of circumcircle of \triangle ABC equals 3 cm.
 - (a) 6

- (b) 9
- (c) 12
- (d) 24
- If $f(x) = \frac{1}{x-2}$, then the graph that represents the function f is









28 In the opposite figure:

ABCD is a parallelogram

$$m (\angle ABD) = 80^{\circ} BD = 7 cm.$$

AB = 5 cm., then the perimeter

of parallelogram = to the nearest cm.



(b) 26

(c) 29

(d) 30

Second

Essay questions

Answer the following questions:

- 1 Find: $\lim_{X \to -2} \frac{3 X^2 12}{X + 2}$
- 2 Graph the function $f: f(x) = \begin{cases} -x^3, & x < 0 \\ x, & x \ge 0 \end{cases}$, from the graph find the range and its type whether it is odd, even or otherwise, and discuss its monotony.
- If $f(x) = 2^x$, find the value of x which satisfies: f(x+1) f(x-1) = 24
- $\lim_{x \to \infty} \frac{4x^5 + 5}{8x^5 + x^4 2}$

Model

5

Interactive test 5



First Multiple choice questions

Choose the correct answer from the given ones:

- $\lim_{x \to 1} \frac{x^{6\frac{1}{2}} x^{\frac{1}{2}}}{x^{3\frac{1}{2}} x^{\frac{1}{2}}} = \dots$
 - (a) $\frac{13}{7}$

- (b) 1
- (c) 2

(d) X

- 2 If $5^{X+1} = 7^{X+1}$, then $3^{X+1} = \dots$
 - (a) zero

- (b) 3
- (c) 2

(d) 1

- 3 If x < 1, then $|3 x| |x 4| = \dots$
 - (a) 1

- (b) 1
- (c) 2×-7
- (d) 7 2 X
- The solution set in \mathbb{R} of the equation : $|2 \times -4| = |\times +1|$ equals
 - (a) $\{1\}$

- (b) {5}
- (c) $\{1,5\}$
- (d) Ø

- 5 The domain of the function $f: f(x) = \sqrt{x-2}$ is
 - (a) IR

- (b) $\{2\}$
- (c) $[2, \infty[$ (d) $]2, \infty[$

- 6 $\lim_{x \to \infty} \frac{\sqrt{4x^2 + 1}}{x 2} = \dots$
 - (a) 4

- (b) 5
- (c) $-\frac{5}{2}$
- (d) 2
- In \triangle ABC, if m (\angle A) = 30°, b = 15 $\sqrt{3}$ cm., m (\angle B) = 60° , then $a = \cdots cm$.
 - (a) 30

- (b) 45
- (c) 15
- (d) 60

- B $\lim_{x \to \infty} (3 + 5 x^2 + 3 x) = \dots$
 - (a) does not exist.
- (b) 5
- (c) ∞
- (d) 11
- The domain of the function f: f(X) = 5 is
 - (a) $\left\{\frac{1}{5}\right\}$

- (b) $\{5\}$
- (c) R
- (d) $\mathbb{R} \{5\}$

- 10 If $f(a) = 2^a$, then $\log_2 f(a) = \dots$
 - (a) 2

- (b) f (a)
- (c) a
- (d) $\frac{1}{2a}$

III From the opposite figure :


$$\lim_{x \to 3} f(x) = \dots$$

(a) 1

(b) 3

(c) - 1

(d) does not exist.



- 12 $\lim_{x \to 0} (2 x^2 + 3) = \dots$
 - (a) 2

- (b) 3
- (c) 5

- (d) 7
- From the following functions, the even function is $f: f(X) = \cdots$
 - (a) $\sin x$

- (b) sin 30°
- (c) X cos X
- (d) $\chi^2 + \tan \chi$

- **14** In \triangle XYZ, $2 \times x \times \dots = x^2 + z^2 y^2$
 - (a) cos X
- (b) cos Z
- (c) cos Y
- (d) sin Y

- 15 If $\lim_{x \to -1} \frac{x^2 + kx + m}{x^2 1} = 3$, then $k + m = \dots$

- (d) 9
- **16** The range of the function $f: f(x) = \frac{x^2 1}{x 1}$ is
 - (a) R
- (b) $\mathbb{R} \{0\}$
- (d) $\mathbb{R} \{2\}$


- 17 If $\frac{a+b}{13} = \frac{b+c}{11} = \frac{c+a}{12}$, then $\cos A = \dots$


 - (a) $\frac{1}{5}$ (b) $\frac{5}{7}$


- (d) $\frac{4}{11}$
- 18 The number of possible solutions of the triangle ABC: $m (\angle A) = 47^{\circ}$, a = 4 cm. , b = 6 cm. equals
 - (a) 1
- (b) 2


(c) 3

- (d) zero
- If $f(x) = 2 (x 1)^3$, then the graph that represents the function f is









- **20** The S.S. of the equation : $\log_{\mathcal{X}}(X+6) = 2$ in \mathbb{R} is
 - (a) $\{3, -2\}$ (b) $\{3\}$
- (c) $\{3,1\}$
- (d) {6,1}
- 21 A man deposite L.E. 12000 in a bank that gives yearly interest 13 %
 - , then the sum of money after 10 years approximately equals L.E.
 - (a) 40735
- (b) 38735
- (c) 36049
- (d) 46030
- In \triangle LMN, m (\angle L) = 30°, MN = 7 cm., then the diameter length of the circle passing through its vertices equals
 - (a) 7 cm.
- (b) 3.5 cm.
- (c) 14 cm.
- The solution set of the equation : $2^{\chi^2} = 16$ in \mathbb{R} is
 - (a) {2}
- (b) $\{-2\}$
- (c) $\{2, -2\}$
- (d) $\{4, -4\}$

The curve : $y = 3(x-5)^2 + 7$ under action of translation 3 units in the positive direction of the X-axis and one unit in the negative direction of the y-aixs is the curve

(a)
$$y = 3 (X + 8)^2 + 6$$

(b)
$$y = 3 (x - 8)^2 + 8$$


(c)
$$y = 3(x-8)^2 + 6$$

(d)
$$y = 3(x + 8)^2 - 6$$

25 In the opposite figure :

BC = 7 cm.,
$$m (\angle A) = 120^{\circ}$$
, AB < AC, then AC = cm.





(a) 5

(d) 4

(c) 8

- The simplest form of the expression: $\frac{1}{\log_x x y z} + \frac{1}{\log_x x y z} + \frac{1}{\log_z x y z} = \frac{1}{\log_z x y z}$
 - (a) Z
- (b) y

(c) 1

(d) X

- In any triangle XYZ, XY: YZ = ·····
 - (a) sin X : sin Y
- (b) sin Y: sin Z
- (c) sin Z: sin X
- (d) sin Z: sin Y
- If the curve y = f(X) represents a real function, then its image by translation 5 units vertically downward is $g(X) = \dots$
 - (a) f(X-5)
- (b) f(x+5)
- (c) f(x) + 5
- (d) f(x) 5


Second **Essay questions**

Answer the following questions :

- 1 Showing steps , find the solution set of the equation : $3^{2X-1} - 4 \times 3^{X} + 9 = 0$, where X is a real number.
- If the function $f: f(x) = \frac{1}{x}$, then find the domain of the function f and the coordinates of the symmetric point of the curve of this function , then find in ${\mathbb R}$ the solution set of the equation : $f\left(\frac{1}{x}\right) = 4$
- Find: $\lim_{x \to 0} \frac{\sqrt{x+4}-2}{x}$

4 Study the opposite figure, then find:

- (1) f(1)
- (2) $\lim_{x \to 1} f(x)$



Model

6

Interactive test 6



Multiple choice questions **First**

Choose the correct answer from the given ones:

- If $\lim_{x \to 4} \frac{x^2 + 7x + b}{x^2 6x + 8} = \frac{15}{2}$, then $b = \dots$
 - (a) 44

- (b) 7
- (c) 8
- (d) 8
- The vertex point of the curve of the function $f: f(x) = x^2 + 3$ is
 - (a)(3,0)
- (b)(0,3)
- (c)(-3,0)
- (d)(0,-3)
- 3 If $\log_a (X + 2) \log_a (X 1) = \log_a 4$, then $X = \dots$
 - (a) 2

- (b) 2
- (c) 1

- (d) 1
- 4 All the functions defined by the following rules are odd except
 - (a) $f(X) = \tan X$
- (b) $f(X) = \csc X$ (c) $f(X) = 7 X^3$ (d) $f(X) = \cos X$

- $\lim_{x \to 0} \frac{x^2 1}{x} = \dots$
 - (a) zero

- (b) 1
- (c) does not exist.
- (d) 1

- i If $x^{\frac{7}{2}} = 64$, then x = ...
 - (a) 512

- (b) 16
- (c) 4
- (d)2
- 11 The area of the circle passing through the vertices of the equilateral triangle ABC whose side length is 9 cm. equals cm².
 - (a) 9 T

- (b) $9\sqrt{3}\pi$
- (c) 27π
- (d) 81 T

- 18 If $f(x) = 3^x$, then the solution set in \mathbb{R} of the equation : f(x-2) + f(x-1) = 36
 - (a) $\{9\}$

- (b) $\{4\}$
- $(c) \{2\}$
- $(d) \{3\}$

- 9 Lim $\frac{(X+h)^7 X^7}{h} = \dots$
 - (a) x^7

- (b) 7×6
- (c) zero
- (d) 1

- 10 In \triangle ABC, $a^2 + b^2 c^2 = \dots$
 - (a) cos A

- (b) a b cos C
- (c) cos C
- (d) 2 a b cos C
- 111 The curve g(x) = |x + 3| is the same as the curve f(x) = |x| by translation 3 units in the direction of
 - (a) OX

- (b) Ox
- (c) Oy
- (d) Ov
- 12 The solution set of the inequality: $|3-2x| \le 1$ in \mathbb{R} is
 - (a) [1, 2]
- (b) 1,2
- (c) $\mathbb{R}]1, 2[$ (d) $\mathbb{R} [1, 2]$
- If (2, 3) lies on the curve of an odd function, then the point lies on the curve of the same function.
 - (a) (-2, -3)
- (b) (2, -3) (c) (-2, 3) (d) (3, 2)

- The point of symmetry of the function $f: f(x) = \frac{2x-1}{x}$ is
 - (a) (1, 1)
- (b) (2, 1)
- (c)(1,2)
- (d) (0, 2)

- 15 $\lim_{x \to 1} \frac{2x-4}{x-2} = \dots$
 - (a) 1

- (b) 2
- (c) 2
- (d) zero
- 16 The range of the function $f: f(x) = \frac{5}{x} + 2$ is
 - (a) IR

- (b) $\mathbb{R} \{2\}$ (c) $\{2\}$
- (d) $\mathbb{R} \{0\}$
- The range of the function $f: f(X) = \begin{cases} 0, & X \le 0 \\ 1, & X > 0 \end{cases}$ is
 - (a) $\{1\}$

- (b) $\{0\}$
- (c) R
- $(d) \{0,1\}$

18 The radius length of the circumcircle of the triangle XYZ in which:

x = 3 cm., y = 5 cm., z = 7 cm. approximately equalscm.

(a) 6


- (b) 8
- (c) 4
- (d) 2

19 In the opposite figure:

If the perimeter of \triangle ABC = 42 cm. and the circle M is the inscribed circle in it, then $m (\angle A) = \cdots$

(a) 53° 7

- (b) 67° 23
- (c) 36° 53
- (d) 22° 37



- **20** If $5^{x-3} = 4^{3-x}$, then $x = \dots$
 - (a) $\frac{5}{4}$

- (c) $\frac{4}{5}$
- (d) zero
- The numerical value of the expression $\frac{\log 64}{\log 8}$ equals
 - (a) 2

- (b) 8
- (c) 80
- (d) 72
- In \triangle DEF, m (\angle D) = 80°, m (\angle E) = 60°, if f = 12 cm., then d =cm.
 - (a) $\frac{12 \sin 80^{\circ}}{\sin 40^{\circ}}$

- (b) $\frac{12 \sin 80^{\circ}}{\sin 60^{\circ}}$ (c) $\frac{12 \sin 40^{\circ}}{\sin 80^{\circ}}$ (d) $\frac{12 \cos 80^{\circ}}{\cos 40^{\circ}}$
- $\lim_{X \to \infty} \frac{(2 X + 1) (4 X 1)^2}{(2 X + 3)^3} = \dots$

- (c) 1
- (d) 8

- If $\lim_{x \to 1} \frac{b}{x+1} = 5$, then $b = \cdots$
 - (a) 5

- (b) 1
- (c) 1

- (d) 10
- 25 In \triangle ABC, if $b^2 = (c a)^2 + ca$, then m (\angle B) =
 - (a) 30°

- (b) 60°
- (c) 90°
- (d) 120°
- 26 The absolute inequality that represents mark of a student from 50 to 70 marks is
 - (a) |x-20| < 10

(b) |x-60| < 10

(c) $|x - 60| \le 10$

- (d) $|x-20| \le 10$
- In \triangle ABC, $\cos(A + B) = \cdots$

- (a) $\frac{a^2 + b^2 c^2}{2ab}$ (b) $\frac{a^2 + c^2 b^2}{2ab}$ (c) $\frac{b^2 + c^2 a^2}{2bc}$ (d) $\frac{c^2 a^2 b^2}{2ab}$

28 In the opposite figure:


 \overline{AD} // \overline{BC} , AB = 4 cm., m ($\angle DAC$) = 40° , m ($\angle B$) = 60° , then the length of $AC \simeq \cdots cm$.

(a) 5

(b) 3

(c) 2

(d) 4



Second **Essay questions**

Answer the following questions :

- In Graph the curve of the function f: f(X) = |X + 2| + 1 and deduce its range and discuss its monotoncity and its type whether it is even, odd or otherwise.
- 2 Find: $\lim_{x \to \infty} (\sqrt{x^2 + 5x} x)$
- 3 Without using calculator find the value of :

$$\log_2 \frac{3}{25} + 5 \log_2 5 + \log_2 27 - \log_2 \frac{125}{12} - \log_2 243$$

4 Find:
$$\lim_{x \to 0} \frac{x^2}{3 x^3 - 2 x^2}$$

Model

Interactive test 7



Multiple choice questions First

Choose the correct answer from the given ones:

- 11 The range of the function f: f(X) = |X| is
 - (a) $[0, \infty[$

- (b) $]0, \infty[$ (c) $]-\infty, 0]$ (d) $]-\infty, 0[$
- In \triangle ABC, $\frac{a}{a+b} = \frac{\sin A}{\cos A}$
 - (a) sin B

- (b) sin C
- (c) $\sin A + \sin B$
- $(d) \sin A + \sin C$
- In \triangle ABC, if $\sin A = 2 \sin C$, BC = 6 cm., then AB = cm.
 - (a) 2

- (b) 3
- (c) 4
- (d) 6

- The solution set in \mathbb{R} of the equation : $x^{\frac{4}{3}} 10 x^{\frac{2}{3}} + 9 = 0$
 - (a) $\{1, 27\}$

(b) $\{-1,1\}$

(c) $\{-1, 1, 27\}$

(d) $\{-1, 1, -27, 27\}$

- $\lim_{X \to \frac{\pi}{A}} \frac{\tan X}{X} = \dots$
 - (a) $\frac{\pi}{2}$

- (b) $\frac{4}{\pi}$
- (c) 1

(d) does not exist.

- If $\sqrt[3]{x^2} = 4$, then $x = \dots$
 - (a) 8

- (b) 8
- $(c) \pm 8$
- $(d) \pm 4$

- In \triangle ABC, $b^2 + c^2 a^2 = 2 b c \times \dots$
 - (a) $\sin (90^{\circ} B)$
- (b) $\sin (90^{\circ} A)$
- (c) cos B
- (d) $\cos (90^{\circ} B)$

- B $\lim_{x \to 1} \frac{4 \sqrt{x + 15}}{1 x^2} = \dots$
 - (a) 16

- (b) 16
- (c) $\frac{1}{16}$
- $(d) \frac{-1}{16}$
- If the radius length of the circle passing throught the vertices of \triangle ABC equals 6 cm.
 - , then $\frac{2 \text{ a}}{\sin A} = \cdots \cos \alpha$.
 - (a) 12

- (b) 6
- (c) 18
- (d) 24
- 10 The solution set of the equation: $(\log_5 y)^2 7 \log_5 y + 12 = 0$ in \mathbb{R} is
 - (a) $\{25, 125\}$
- (b) $\{25, 625\}$ (c) $\{\frac{1}{25}, 625\}$ (d) $\{125, 625\}$
- 11 The solution set of the equation : |X| + 3 = 0 in \mathbb{R} is
 - (a) $\{3\}$

- (b) $\{-3\}$
- $(c)\{0\}$
- $(d) \emptyset$

- 12 If $\lim_{x \to 1} \frac{x^2 k^2}{x + 2} = -1$, then $k = \dots$

- (c) 4
- $(d) \pm 2$
- 13 If f is an odd function then $\frac{5 f(x) + 2 f(-x)}{4 f(x)} = \frac{1}{2}$
 - (a) $\frac{7}{4}$

- (d) $\frac{5}{4}$
- In \triangle XYZ, x = 5 cm., y = 7 cm., m (\angle Z) = 65°, then z approximately equalscm.
 - (a) 7.6

- (b) 6.7
- (c)7.8
- (d) 8.7

- 15 If $\log_2 x = 3$, then $\log_x 2 = \dots$
 - (a) 2


- (b) $\frac{1}{3}$
- (c) 8

(d) 9

16 The rule of the function represented in the opposite

figure is $f(X) = \cdots$

- (a) $(x-2)^2 + 1$
- (b) $-(x-2)^2+1$
- $(c) (x-1)^2 + 2$
- (d) $(-x+1)^2+2$



- 11 The solution set of the equation : $|2 \times -1| = 5$ in \mathbb{R} is
 - (a) $\{3\}$

- (c) Ø
- (d) $\{3, -2\}$
- 18 If $f(x) = \begin{cases} x 4 & , & x \ge 4 \\ g(x) & , & x < 4 \end{cases}$ is symmetric about the straight line X = 4

, then the function g is

(a) an increasing function.

(b) a decreasing function.

(c) an even function.

(d) a constant function.

- $\lim_{x \to 0} \frac{3 x + 2 x^{-1}}{x + 4 x^{-1}} =$
 - (a) $\frac{1}{4}$

- (b) $\frac{1}{2}$
- (c) 2
- (d) 4
- The domain of the function $f: f(x) = \frac{x+2}{x^2-9}$ is
 - (a) $\{3, -3\}$
- (b) $\mathbb{R} \{3, -3\}$ (c) $\mathbb{R} \{3\}$
- (d) R

- $\lim_{x \to \infty} \left(\frac{1}{x-2} + 1 \right) = \dots$

- (c) zero
- (d) ∞
- The solution set in \mathbb{R} of the equation : $\sqrt{x^2 6x + 9} = 9$ is
 - (a) $\{-6, 12\}$
- (b) {12}
- (c) $\{-6\}$ (d) $\{6, -12\}$
- If the curve of the function $f: f(X) = \log_4 (1 aX)$ passes through $\left(\frac{1}{8}, -\frac{1}{2}\right)$, then a =
 - (a) 3

- (b) 2
- (c) 4

(d) 8

In \triangle ABC, if b = c, then $\cos C = \cdots$

(a)
$$\frac{a}{2b}$$

$$(b) \frac{b}{2c} \qquad (c) \frac{2b}{c}$$

(c)
$$\frac{2b}{c}$$

(d)
$$\frac{2b}{a}$$

25 The solution set of the inequality $|2 \times + 3| \le 7$ in \mathbb{R} is

(a)
$$]-5,2[$$

(b)
$$]-2,5[$$
 (c) $[-2,5]$

(c)
$$[-2,5]$$

(d)
$$[-5, 2]$$

 $\lim_{x \to \frac{\pi}{2}} (1 - \cos x + \sin x) = \cdots$

$$(a) - 1$$

27 \triangle ABC in which a = 4 cm., $b = 4\sqrt{3}$ cm., c = 8 cm., then sine of the smallest angle measure in it = ······

(a)
$$\frac{1}{2}$$

(b)
$$\frac{\sqrt{3}}{2}$$

28 In \triangle ABC, if m (\angle C) = 60°, $a^2 + b^2 - c^2 = k \ a \ b$, then $k = \dots$

(a)
$$\frac{1}{2}$$

$$(d) - 1$$

Essay questions Second

Answer the following questions:

1 Without using the calculator, find the value of:

$$\log_2 \frac{3}{25} + 5 \log_2 5 + \log_2 27 - \log_2 \frac{125}{12} - \log_2 243$$

2 Find: $\lim_{x \to \infty} \frac{2x-3}{\sqrt[3]{125 x^3 + 5}}$

Graph the function $f: f(X) = \begin{cases} x^2, & x < 0 \\ x, & x \ge 0 \end{cases}$ and determine the range and monotonicity.

 $\lim_{x \to 0} \frac{(x+2)^2 - 4}{x^2 + x}$

Model

Interactive test 8



First Multiple choice questions

Choose the correct answer from the given ones:

- If $\log_2 X = 3$, then $X = \dots$
 - (a) 6

- (b) 2
- (c) 9
- (d) 8

- $\lim_{x \to 0} \sqrt{64 + x^2} = \dots$
 - (a) 64

- (b) 16
- (c) 8

- (d) otherwise.
- The diameter length of the circle inscribed in an equilateral triangle whose side length is $4\sqrt{3}$ cm. equals cm.
 - (a) $2\sqrt{3}$

- (b) 4 \(\frac{1}{3}\)
- (c) 4
- (d) 8
- If y = f(X) is a real function, then its image by translation 2 units right is $g(X) = \cdots$
 - (a) f(X-2)
- (b) f(X+2)
- (c) f(x) + 2
- (d) f(x) 2
- The number of possible solutions of \triangle ABC where m (\angle A) = 60°, b = 3 cm. , a = 5 cm. is
 - (a) 1

- (b) 2
- (c) no solution.
- (d) an infinite number of triangles.
- $\lim_{X \longrightarrow \text{zero}} \frac{X^2 + X}{X} = \dots$
 - (a) zero

- (b) 1
- (c) 2
- (d)3
- If $f(X) = 5^X$, then the solution set in \mathbb{R} of the equation : f(X) + f(X 1) = 150equals
 - (a) $\{3\}$

- (b) $\{5\}$
- (c) $\{2\}$
- (d) $\{3,5\}$

- In \triangle ABC, $\cos(A + B) = \cdots$
 - (a) $\frac{a^2 + b^2 c^2}{2 a b}$

- (b) $\frac{a^2 + c^2 b^2}{2 a c}$ (c) $\frac{b^2 + c^2 a^2}{2 b c}$ (d) $\frac{c^2 a^2 b^2}{2 a b}$
- - (a) (1,0)
- (b) (-1,0) (c) (0,1)
- (d)(0,-1)

- $\lim_{x \to 3} \frac{x^2 7x + 12}{x 3} = \dots$

- (b) 1
- (c) 2
- (d) 7
- 11 The domain of the function $f: f(x) = \frac{3}{\sqrt{x+4}}$ equals
 - (a) [-4,∞[
- (b) $]-\infty,4]$
- (c)]-4,∞[
- (d) $]-\infty, -4[$

12 In the opposite figure:


If AC = 20 cm.

- , then the perimeter of \triangle ACM \simeq cm.
- (a) 41.6

(b) 43.5

(c) 45

(d) 47.5



- $\log 2 + \log 5 = \dots$
 - (a) 1

- (b) log 7
- (c) 10
- (d) log 5
- 14 The domain of the function $f: f(x) = \sqrt{9-x}$ is
 - (a) R

- (b) $\mathbb{R} \{9\}$ (c) $]-\infty, 9]$
- (d) [9,∞[
- 15 In \triangle ABC, c = 19 cm., $m (\angle A) = 112^{\circ}$, $m (\angle B) = 33^{\circ}$, then the area of \triangle ABC to the nearest cm² equals cm².
 - (a) 64

- (b) 128
- (c) 185
- (d) 159
- 16 The solution set of the inequality : |X| 1 > 0 in \mathbb{R} is
 - (a) $\mathbb{R} [-1, 1]$
- (b)]-1,1[(c) $\mathbb{R}-]-1,1[$ (d) [-1,1]

- $\lim_{x \to -2} \left| \frac{1}{x} \right| = \dots$
 - (a) 1

- (b) 1
- $(c) \frac{1}{2}$
- (d) $\frac{1}{2}$

- $\lim_{x \to \infty} \frac{1}{x^2 3x + 2} = \dots$
 - (a) 3

- (b)9
- (c) 27
- (d) 81

- 19 Which of the following does not equal $(\sqrt[5]{x^4})$?
 - (a) $(\sqrt[5]{x})^4$
- (c) $\chi \frac{4}{5}$
- (d) $\left(\chi^{\frac{1}{5}}\right)^4$

- If the function f is even in [c,d], then $c+d=\cdots$
 - (a) 2 c

- (b) 2 d
- (c) c d
- (d) zero

- 21 If $\sqrt[3]{x^2} = 9$, then $x \in \dots$
 - (a) {27}

- (b) $\{27, -27\}$ (c) $\{1\}$
- (d) Ø
- If $\left(\frac{1}{2}\right)^{a^2-a-2} = 1$, where a > zero, then $a = \cdots$
 - (a) 1

- (b) 3
- (c) 2

- (d) 3
- Which of the functions defined by the following rules represents an exponential function increasing on its domain \mathbb{R} ?
 - (a) $y = 3 (1.05)^{x}$
- (b) $y = \frac{1}{3} \left(\frac{1}{1.5} \right)^X$ (c) $y = 3 + (0.5)^X$ (d) $y = (0.5)^X$
- In \triangle ABC, if $2 \sin A = 3 \sin B = 4 \sin C$, then $a : b : c = \cdots$
 - (a) 2:3:4

- (b) 4:3:2 (c) 3:4:6 (d) 6:4:3
- If $\lim_{x \to a} \frac{a x}{3} = 12$, then $a = \dots$
 - $(a) \pm 12$

- $(b) \pm 6$
- (c) 4
- (d) $\frac{1}{6}$

- **26** If |X| + |X 3| = 3, then X(X 3)zero
 - (a) <

- (b) >
- (c) ≤
- (d) ≥

- In $\triangle XYZ$, if X = y, then $\cos X = \cdots$
 - (a) $\frac{2 y^2}{}$

- (b) $\frac{z}{2v}$
- (c) $\frac{z}{4x}$
- $(d) \frac{y}{2x}$
- The perimeter of \triangle ABC, in which b = 11 cm., $m (\angle A) = 67^{\circ}$, $m (\angle C) = 46^{\circ}$ equals (to the nearest cm.)
 - (a) 22

- (b) 38
- (c) 31
- (d) 27

Second **Essay questions**

Answer the following questions :

Without using the calculator find the value of:

$$\log_3 54 - \log_3 \frac{8}{15} + \log_3 \frac{4}{5}$$

- 2 Find: $\lim_{x \to 5} \frac{x^2 5x}{\sqrt{x + 4} 3}$
- Graph the curve of the function $f: f(x) = (x+2)^3 + 1$ and from the graph deduce the range and its monotony and its type whether it is even \cdot odd or otherwise.
- $\lim_{x \to \infty} \frac{5 x^{-3} + 4 x^{-2} 3}{7 x^{-3} 2 x^{-2} + 8}$

Model

9

Interactive test 9



First Multiple choice questions

Choose the correct answer from the given ones:

- 11 The solution set of the equation: $\log_{(X+3)} 125 = 3$ in \mathbb{R} is
 - (a) $\{5\}$
- (b) {3}
- (c) Ø
- (d) $\{2\}$
- 2 Δ LMN in which m (\angle L) = 30°, m = 9 cm. has two solutions when ℓ = cm.
 - (a) 6


- (b) 10
- (c) 11

- (d) 2
- If $4 = \log_2 x$, then the equivalent exponential form is
 - (a) $X^2 = 4$
- (b) $X^4 = 2$
- (c) $x = 2^4$
- (d) $2^{x} = 4$

The domain of the function represented by the opposite figure is



$$(c)] - \infty , 0$$



- 5 If $f: \mathbb{R} \longrightarrow \mathbb{R}$ where f(X+1) f(X) = X 1, then $f(10) f(9) = \cdots$
 - (a) 1

- (b) 9
- (c) 8

(d) 18

- 6 $\lim_{x \to 0} \frac{x^2 + x}{x^3 + x} = \dots$
 - (a) $\frac{2}{3}$

- (b) 1
- (c) zero
- (d) does not exist.

- The image of the curve f(x) = |x| 5 by translation 3 units in the direction of Ox and 5 units in the direction of Oy is
 - (a) g(X) = |X 3| + 5

(b) g (X) = |X - 3|

(c) g(X) = |X - 3| - 10

- (d) g (X) = |X + 3|
- $\lim_{X \to \infty} \frac{\sqrt{4 x^2 + 7 + 3 X}}{2 x + 9} = \dots$

- (b) $\frac{5}{2}$
- (c) $\frac{5}{4}$

- (d) $\frac{5}{9}$
- - (a)]-1,3[
- (b) $\mathbb{R} [-1, 3]$ (c)]-2, 2[
- $(d) \emptyset$

- In \triangle ABC, c (a cos B + b cos A) =
 - (a) a²

- (b) b^2

- (d) $2 c^2$
- 11 ABCD is a parallelogram in which: AB = 9 cm., BC = 13 cm., AC = 20 cm., then the length of BD equals cm.
 - (a) 10

- (b) 5
- (c) 18.5
- (d) 20
- 12 If the domain of the function $f: f(x) = \frac{2}{x^2 6x + k}$ is $\mathbb{R} \{3\}$, then $k = \dots$
 - (a) 3

 $(d) \pm 9$

- $\lim_{X \to 4} \frac{X^3 64}{X 4} = \dots$
 - (a) 96

- (b) 48
- (c) 32

- (d) 16
- If $f(x) = \frac{\sqrt{x^2 2x + 1}}{x 1}$, then the range of the function f is
 - (a) {1}

- (c) [-1,1]
- (d) $\{-1,1\}$
- The solution set of the following equation in \mathbb{R} : $\log_2 x \frac{3}{\log_2 x} = 2$ equals
 - (a) $\{\frac{1}{2}\}$

- (b) $\{8,2\}$
- (c) $\left\{8, \frac{1}{2}\right\}$
- (d) $\{2\}$

- $\lim_{h \to 0} \frac{(x+h)^9 x^9}{h} = \dots$

- (b) $9x^8$
- (c) zero
- (d) does not exist.

- $\log_3 15 \log_3 5 = \dots$
 - (a) 3

- (b) 1
- (c) zero
- (d) 3

- 18 If ABC is a triangle in which a = 4 cm., $b = 4\sqrt{3}$ cm., c = 8 cm., then sine of its smallest angle equals
 - (a) $\frac{1}{2}$


- (b) $\frac{\sqrt{3}}{2}$
- (c) 1

(d) zero

- If $x = 5 + 2\sqrt{6}$, then $\log\left(\frac{1}{x} + x\right) = \dots$
 - (a) 1

- (b) $5 2\sqrt{6}$
- (c) 10

- (d) $5 + 2\sqrt{6}$
- 20 Which of the functions represented graphically as follows is neither even nor odd?



21 In the opposite figure :


If $tan (\angle DEC) = \frac{3}{4}$, then the radius length of the circumcircle of \triangle ABC = cm.

(a) 9

(b) 5.7

(c) $4\frac{3}{4}$

(d) 3.75



- The solution set of the equation: $\frac{1}{\log_2 x} + \frac{1}{\log_3 x} = 2$ is
 - (a) $\{\sqrt{6}\}$
- (b) $\{-\sqrt{6}\}$ (c) $\{\sqrt{6}, -\sqrt{6}\}$ (d) $\{6\}$


- **23** If $\sqrt[3]{x^2} = 9$, then $x \in \dots$
 - (a) $\{-81, 81\}$ (b) $\{-27, 27\}$ (c) $\{-9, 9\}$ (d) [3, 7]

In the opposite figure :

 $f(x) = \cdots$

(a) $\frac{1}{x-1}$

- (b) $\frac{1}{|x-1|}$
- (c) $|x^2 1|$
- (d) $|x-1|^2$



$$\lim_{x \to \infty} \left(x - \sqrt{x^2 + 5x} \right) = \dots$$

(a) 2

- (b) 3
- (c) $\frac{-5}{2}$
- (d) $\frac{1}{4}$

In
$$\triangle XYZ$$
, if $\sin X = 2 \sin Z$, $YZ = 6 \text{ cm.}$, then the length of $\overline{XY} = \cdots \text{ cm.}$

(a) 12

- (b) 2
- (c) 6

(d) 3

(a) $7\sqrt{3}$

- (b) $14\sqrt{3}$
- (c) 7

(d) 14

 $(a) - (\cos B + \cos C)$

(b) cos B - cos C

 $(c) \cos (B + C)$

 $(d) - \cos (B + C)$

Second Essay questions


Answer the following questions:

- 1 Graph the function $f: f(x) = \sqrt{x^2 4x + 4}$ and determine its range and discuss its monotony.
- Graph the curve of the function $f: f(x) = x^3 5$ and from the graph discuss the monotonicity of the function and show its type whether it is even, odd or otherwise.

3 Find:
$$\lim_{x \to 0} \frac{\sqrt{9x+16}-4}{x}$$

4 Study the opposite figure, then find:

- (1) f(0)
- (2) $\lim_{x \to 0} f(x)$
- (3) f(2)
- $(4) \lim_{X \to 2} f(X)$



Model 10

Interactive test 10



First Multiple choice questions

Choose the correct answer from the given ones:

- 1 The solution set of the equation $\log_3 (X-4) + \log_3 (X+4) = 2$ in \mathbb{R} is
 - (a) $\{5\}$

- (b) $\{5, -5\}$ (c) $\{3, -3\}$ (d) $\{3, 5\}$

- $\lim_{x \to 0} \frac{(x+2)^2 4}{x^2 + x} = \dots$
 - (a) zero

- (b) 2
- (c) 4

- (d) 8
- 3 If the ratio among the measures of the angles of a triangle is 8:3:1, then the ratio between the longest two sides in the triangle is
 - (a) $\sqrt{3}:2$
- (b) $\sqrt{6}:2$
- (c) 8:3
- (d) 8:5

- $\lim_{X \to -3} \frac{\sqrt{x+7} 2}{x+3} = \dots$
- (b) $\frac{1}{2}$
- (c) 2

(d) 4

- 5 If $3^a = 4^b$, then: $9^{\frac{a}{b}} + 16^{\frac{b}{a}} = \dots$
 - (a) 7

- (c) 20

(d) 25

- If $\lim_{x \to \infty} \frac{3 k |x|}{4 x + 3} = 6$, then $k = \dots$

- (b) $\frac{3}{4}$
- (c) 8

- (d) 3
- If $f(x) = x^3$, then the image of the curve of f by reflection in X-axis and translation 3 units in the direction of \overrightarrow{OX} and two units in the direction of \overrightarrow{OY} is
 - $(a) (x-3)^3 2$

(b) $-(x+3)^3+2$

 $(c) - (x+3)^3 - 2$

- $(d) [(x+3)^3 + 2]$
- B If $2^{x-3} = 1$, then $x = \dots$
 - (a) 3

- (b) 3
- (c) 1

- (d) zero
- If $a \in \mathbb{R}^+ \{1\}$, x, $y \in \mathbb{R}^+$, $\log_a y \neq 0$, then $\frac{\log_a x}{\log_a y} = \cdots$
 - (a) $\log_a \frac{x}{y}$
- (b) $\log_a (X y)$ (c) $\log_a X \log_a y$ (d) $\log_y X$

Final examinations

$$\frac{1}{\log_2 30} + \frac{1}{\log_3 30} + \frac{1}{\log_5 30} = \dots$$

(a) 1

(b) log₆ 5

(c) log 30

(d) 30

In \triangle ABC, m (\angle A) = 112°, m (\angle B) = 33°, c = 19 cm. , then b to the nearest cm. = cm.

(a) 16

(b) 17

(c) 18

(d) 20

12 If $2^{x} = 20$, n < x < n + 1, n is an integer, then $n = \dots$

(a) 4

(b) 5

(c) 6

(d) 10

(a) cos X

(b) sin Z

(c) cos Z

(d) sin X

 $\lim_{x \to 1} \frac{x^2 + 5x - 6}{x^2 - 1} = \dots$

(a) 1

(b) 5

(c) 6

(d) 3.5

15 The exponential function whose base is a , is increasing if

(a) a > 0

(b) a > 1

(c) 0 < a < 1

(d) a = 1

16 $\lim_{x \to 2} \frac{x^5 - 32}{x^4 - 16} = \dots$

(a) 2

(b) 20

(c) $\frac{5}{4}$

11 If f is an odd function, a \subseteq the domain of f, then $f(a) + f(-a) = \cdots$

(a) 2 f (a)

(b) 2 f (-a)

(c) zero

(d) f (a)

111 The solution set in \mathbb{R} of the equation : |x-3| = |9-2|x| equals

(a) $\{4\}$

(b) {4,6} (c) {6}

(d) {2,6}

The range of the function $f: f(X) = \begin{cases} 2 X + 3 &, & X > 3 \\ 9 &, & X < 3 \end{cases}$ is

(a) $\{3\}$

(b) R

(c) 9,∞

(d) [9,∞[

Diameter length of the circumcircle of equilateral triangle whose side length $10\sqrt{3}$ cm. equals cm.

(a) 5

(b) 10

(c) 15

(d) 20

 $\lim_{X \to 0} \frac{(X+1)^{12} - 1}{X} = \dots$

(b) 6

(c) zero

(d) 12

If the area of \triangle ABC is "X" and the radius length of its circumcircle is "r"

, then
$$\frac{4 \text{ r } \chi}{\text{a b c}} = \dots$$

 $\frac{a}{\sin A}$

- (b) cos A
- (c) 1

- (d) r
- If $f(x) = 7^{x+1}$, then the value of x which satisfies: f(2x-1) + f(x-2) = 50equals
 - (a) 1

- (b)7
- (c) zero
- (d) 2
- If L, M are the roots of the equation: $x^2 4x + 4 = 0$, then $\log_2 L + \log_2 M = \dots$
 - (a) 2

- (b) 2
- (c) 4
- (d) 4

- 25 If $\sqrt{x^2 2x + 1} > 4$, then $x \in \dots$
 - (a) [-3,5]

- (b)]-3,5[(c) $\mathbb{R}-]-3,5[$ (d) $\mathbb{R}-[-3,5]$
- In triangle ABC, a = 4 cm., b = 7 cm., $m (\angle A) = 112^{\circ}$, then the number of triangles satisfy these conditions equals
 - (a) 1

- (b) 2
- (c) 0

- (d) infinite number.
- In triangle ABC, $m (\angle A) : m (\angle B) : m (\angle C) = 2 : 3 : 4$, AB = 12 cm., then the length of AC = cm.
 - (a) 10

- (b) 11
- (c) 16
- (d) 18

- **28** In \triangle ABC: $\frac{c^2 a^2 b^2}{2ab} = \cdots$
 - $(a) \cos (A + B)$
- (b) cos C
- (c) $\sin (C + 90^{\circ})$
- $(d) \cos (B + C)$

Second **Essay questions**

Answer the following questions :

- Find: $\lim_{x \to 2} \frac{(x-1)^6 1}{x-2}$
- Without using the calculator prove that : $\log_5 \frac{15}{7} + \log_5 \frac{35}{3} \log_5 \frac{1}{5} = \log_2 8$
- Betermine the type of the function $f: f(X) = X^2 + \sin X$ whether it is even, odd or otherwise.
- $\lim_{x \to \infty} (x^3 + 5x^2 + 1)$



ကြောင်္ကျာပိုက်ကြောင်္ကြာကြောင်းကြောင်ကြောင်းကြောင်



وثلاراي لطبع العثمات من عثمت 4 الباعثمان والباعثمان وال

